

Mouse Oxidized Low Density Lipoprotein (OxLDL) ELISA Kit | OxLDL elisa kit

Mouse Oxidized Low Density Lipoprotein (OxLDL) Wide-range ELISA Kit


CATALOG # AAA189168	REACTIVITY Mouse	APPLICATIONS ELISA
-------------------------------	----------------------------	------------------------------

PRODUCT SYNONYMS

Oxidized Low Density Lipoprotein (OxLDL); Mouse Oxidized Low Density Lipoprotein (OxLDL) Wide-range ELISA Kit; Oxidized Low Density Lipoprotein; OL; OxLDL elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Mouse
Specificity	This assay has high sensitivity and excellent specificity for detection of Wide-range Oxidized Low Density Lipoprotein (OxLDL). No significant cross-reactivity or interference between Wide-range Oxidized Low Density Lipoprotein (OxLDL) and analogues was observed.
Assay Type	Double-antibody Sandwich
Samples	Serum, plasma and other biological fluids
Detection Range	3.12-200ng/mL
Sensitivity	The minimum detectable dose of this kit is typically less than 1.34ng/mL
Applicable Applications for OxLDL elisa kit	ELISA
Precision	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level Wide-range Oxidized Low Density Lipoprotein (OxLDL) were tested 20 times on one plate, respectively. Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level Wide-range Oxidized Low Density Lipoprotein (OxLDL) were tested on 3 different plates, 8 replicates in each plate. CV(%) = SD/meanX100 Intra-Assay: CV<10% Inter-Assay: CV<12%
Test Principle	The test principle applied in this kit is Sandwich enzyme immunoassay. The microtiter plate provided in this kit has been pre-coated with an antibody specific to Wide-range Oxidized Low Density Lipoprotein (OxLDL). Standards or samples are then added to the appropriate microtiter plate wells with a biotin-conjugated antibody specific to Wide-range Oxidized Low Density Lipoprotein (OxLDL). Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain Wide-range Oxidized Low Density Lipoprotein (OxLDL), biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm +/- 10nm. The concentration of Wide-range Oxidized Low Density Lipoprotein (OxLDL) in the samples is then determined by comparing the O.D. of the samples to the standard curve.
Preparation and Storage	The stability of kit is determined by the loss rate of activity. The loss rate of this kit is less than 5% within the expiration date under appropriate storage condition. To minimize extra influence on the performance, operation procedures and lab conditions, especially room temperature, air humidity, incubator temperature should be strictly controlled. It is also strongly suggested that the whole assay is performed by the same operator from the beginning to the end.



Standard Curve (Sample)

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/mouse-elisa-kits/oxidized-low-density-lipoprotein-oxldl/189168>