

Mouse Histidine Decarboxylase (HDC) ELISA Kit | HDC elisa kit

Mouse Histidine Decarboxylase (HDC) ELISA Kit

CATALOG #
AAA189858

REACTIVITY
Mouse


PRODUCT SYNONYMS

Histidine Decarboxylase (HDC); Mouse Histidine Decarboxylase (HDC) ELISA Kit; HDC/Histidine Decarboxylase; HDC elisa kit

PRODUCT SPECIFICATIONS

Reactivity	Mouse
Specificity	This assay has high sensitivity and excellent specificity for detection of HDC. No significant cross-reactivity or interference between HDC and analogues was observed.
Assay Type	Quantitative Sandwich
Samples	Mouse serum, plasma, tissue homogenates, cell lysates, cell culture supernates and other biological fluids
Detection Range	0.156-10ng/mL
Sensitivity	< 0.057ng/mL
Intra-assay Precision	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level HDC were tested 20 times on one plate, respectively. Intra-Assay: CV<10%
Inter-assay Precision	Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level HDC were tested on 3 different plates, 8 replicates in each plate. CV(%) = SD/meanX100. Inter-Assay: CV<12%
Preparation and Storage	Store at 4 degree C for 1 month/-20 degree C for long-term (within shelf life) Shelf Life: 8 months The stability of kit is determined by the loss rate of activity. The loss rate of this kit is less than 5% within the expiration date under appropriate storage condition. To minimize extra influence on the performance, operation procedures and lab conditions, especially room temperature, air humidity, incubator temperature should be strictly controlled. It is also strongly suggested that the whole assay is performed by the same operator from the beginning to the end.

IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

Related Product Information for HDC elisa kit	Principle of the Assay: The microplate provided in this kit has been pre-coated with an antibody specific to HDC. Standards or samples are then added to the appropriate microplate wells with a biotin-conjugated antibody specific to HDC. Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain HDC, biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm ± 10nm. The concentration of HDC in the samples is then determined by comparing the O.D. of the samples to the standard curve. Intended Uses: The kit is a sandwich enzyme immunoassay for in vitro quantitative measurement of HDC in mouse serum, plasma, tissue homogenates, cell lysates, cell culture supernates and other biological fluids.
Product Categories/Family for HDC elisa kit	Enzyme & Kinase

NCBI AND UNIPROT INFORMATION

NCBI GI #	341940427
NCBI GeneID	15186
UniProt Accession #	P23738; Q9QWU3
NCBI Official Full Name	Histidine decarboxylase
NCBI Official Synonym Full Names	histidine decarboxylase
NCBI Official Symbol	Hdc
NCBI Official Synonym Symbols	Hdc-a; Hdc-c; Hdc-e; Hdc-s; AW108189
NCBI Protein Information	histidine decarboxylase; L-histidine decarboxylase
UniProt Protein Name	Histidine decarboxylase
UniProt Gene Name	Hdc
UniProt Synonym Gene Names	HDC
UniProt Entry Name	DCHS_MOUSE

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/mouse-elisa-kits/histidine-decarboxylase-hdc/189858>