

Pig Complement Component 3a (C3a) CLIA Kit | C3a clia kit

Pig Complement Component 3a (C3a) CLIA Kit

CATALOG # AAA190102	REACTIVITY Pig
-------------------------------	--------------------------

PRODUCT SYNONYMS

Complement Component 3a (C3a); Pig Complement Component 3a (C3a) CLIA Kit; C3a CLIA kit (Pig); Complement Component 3a; C3a clia kit

PRODUCT SPECIFICATIONS


Reactivity	Pig
Specificity	This assay has high sensitivity and excellent specificity for detection of Complement Component 3a (C3a). No significant cross-reactivity or interference between Complement Component 3a (C3a) and analogues was observed. Reactive with Pig C3a/Complement Component 3a
Assay Type	Double-antibody Sandwich
Samples	Serum, Plasma and other biological fluids
Detection Range	31.2-2,000pg/mL
Sensitivity	11.2pg/mL
Sample Size	100uL
Detection Method	Chemiluminescence
Featured Series	SE kit
Precision	Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level Complement Component 3a (C3a) were tested 20 times on one plate, respectively. Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level Complement Component 3a (C3a) were tested on 3 different plates, 8 replicates in each plate. CV(%) = SD/meanX100 Intra-Assay: CV<10%, Inter-Assay: CV<12%
Function	Detects protein (regular version)
Assay Procedure Summary	1. Prepare all reagents, samples and standards 2. Add 100ul standard or sample to each well. Incubate 2 hours at 37 degree C 3. Aspirate and add 100ul prepared Detection Reagent A. Incubate 1 hour at 37 degree C 4. Aspirate and wash 3 times 5. Add 100ul prepared Detection Reagent B. Incubate 30 minutes at 37 degree C 6. Aspirate and wash 5 times 7. Add 100ul Substrate Solution. Incubate 10 minutes at 37 degree C 8. Read RLU value immediately.
Test Principle	The microplate provided in this kit has been pre-coated with an antibody specific to Complement Component 3a (C3a). Standards or samples are then added to the appropriate microplate wells with a biotin-conjugated antibody specific to Complement Component 3a (C3a). Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. Then the mixture of substrate A and B is added to generate glow light emission kinetics. Upon plate development, the intensity of the emitted light is proportional to the Complement Component 3a (C3a) level in the sample or standard.;
Research Area	Infection immunity, Immune molecule, Autoimmunity

Preparation and Storage

Store at 4 degree C for 1 month/-20 degree C for long-term (One year within shelf life)

The stability of kit is determined by the loss rate of activity. The loss rate of this kit is less than 5% within the expiration date under appropriate storage condition. To minimize extra influence on the performance, operation procedures and lab conditions, especially room temperature, air humidity, incubator temperature should be strictly controlled. It is also strongly suggested that the whole assay is performed by the same operator from the beginning to the end.

IMAGES



Standard Curve (Sample)

NCBI AND UNIPROT INFORMATION

NCBI GI #	47522844
NCBI GeneID	397072
NCBI Accession #	NP_999174.1
NCBI GenBank Nucleotide #	NM_214009.1
UniProt Accession #	P01025; O97940; Q9GKP1
Molecular Weight	186,807 Da
NCBI Official Full Name	complement C3
NCBI Official Symbol	C3
NCBI Official Synonym Symbols	C3d
NCBI Protein Information	complement C3
UniProt Protein Name	Complement C3
UniProt Gene Name	C3
UniProt Synonym Gene Names	C3bc; ASP

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/pig-clia-kits/complement-component-3a-c3a/190102>