

Rabbit L1CAM Polyclonal Antibody | anti-L1CAM antibody

Anti-L1CAM antibody


CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA19138	Rabbit	Human, Mouse, Rat No cross reactivity with other proteins.	Immunohistochemistry

PRODUCT SYNONYMS

L1CAM; Polyclonal Antibody; Anti-L1CAM antibody; Neural cell adhesion molecule L1; N-CAM-L1; NCAM-L1; CD171; CAML1; MIC5; anti-L1CAM antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat No cross reactivity with other proteins.
Clonality	Polyclonal
Form/Format	Lyophilized
Sequence Length	1,248
Applicable Applications for anti-L1CAM antibody	IHC (Immunohistochemistry)
Application Notes	Immunohistochemistry (Paraffin-embedded Section): 0.5-1µg/ml
Immunogen	A synthetic peptide corresponding to a sequence of human L1CAM (NMVITWKPLRWMDWNAPQVQYRVQWRPQGTRGPW).
Subcellular Localization	Cell membrane.
Reconstitution	Add 0.2ml of distilled water will yield a concentration of 500µg/ml.
Contents	Each vial contains 4mg Trehalose, 0.9mg NaCl, 0.2mg Na ₂ HPO ₄ , 0.05mg Na ₃ .
Preparation and Storage	Store at -20 degree C for one year. After reconstitution, at 4 degree C for one month. It can also be aliquotted and stored frozen at -20 degree C for a longer time. Avoid repeated freezing and thawing.




IHC (Immunohistochemistry)

Figure 7. IHC analysis of L1CAM using anti-L1CAM antibody (AAA19138).L1CAM was detected in paraffin-embedded section of human mammary cancer tissue. Heat mediated antigen retrieval was performed in citrate buffer (pH6, epitope retrieval solution) for 20 mins. The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1ug/ml rabbit anti-L1CAM Antibody (AAA19138) overnight at 4 degree C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37 degree C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC) with DAB as the chromogen.




IHC (Immunohistochemistry)

Figure 6. IHC analysis of L1CAM using anti-L1CAM antibody (AAA19138).L1CAM was detected in paraffin-embedded section of human lung cancer tissue. Heat mediated antigen retrieval was performed in citrate buffer (pH6, epitope retrieval solution) for 20 mins. The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1ug/ml rabbit anti-L1CAM Antibody (AAA19138) overnight at 4 degree C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37 degree C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC) with DAB as the chromogen.





IHC (Immunohistochemistry)

Figure 5. IHC analysis of L1CAM using anti-L1CAM antibody (AAA19138).L1CAM was detected in paraffin-embedded section of human glioma tissue. Heat mediated antigen retrieval was performed in citrate buffer (pH6, epitope retrieval solution) for 20 mins. The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1ug/ml rabbit anti-L1CAM Antibody (AAA19138) overnight at 4 degree C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37 degree C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC) with DAB as the chromogen.



IHC (Immunohistochemistry)

Figure 4. IHC analysis of L1CAM using anti-L1CAM antibody (AAA19138).L1CAM was detected in paraffin-embedded section of human colon cancer tissue. Heat mediated antigen retrieval was performed in citrate buffer (pH6, epitope retrieval solution) for 20 mins. The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1ug/ml rabbit anti-L1CAM Antibody (AAA19138) overnight at 4 degree C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37 degree C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC) with DAB as the chromogen.



IHC (Immunohistochemistry)

Figure 3. IHC analysis of L1CAM using anti-L1CAM antibody (AAA19138). L1CAM was detected in paraffin-embedded section of rat brain tissue. Heat mediated antigen retrieval was performed in citrate buffer (pH6, epitope retrieval solution) for 20 mins. The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1ug/ml rabbit anti-L1CAM Antibody (AAA19138) overnight at 4 degree C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37 degree C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC) with DAB as the chromogen.

IHC (Immunohistochemistry)

Figure 2. IHC analysis of L1CAM using anti-L1CAM antibody (AAA19138). L1CAM was detected in paraffin-embedded section of mouse brain tissue. Heat mediated antigen retrieval was performed in citrate buffer (pH6, epitope retrieval solution) for 20 mins. The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1ug/ml rabbit anti-L1CAM Antibody (AAA19138) overnight at 4 degree C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37 degree C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC) with DAB as the chromogen.

ADDITIONAL INFORMATION

Related Product Information for anti-L1CAM antibody

Description: L1, also known as L1CAM, is a transmembrane protein member of the L1 protein family, encoded by the L1CAM gene. The protein encoded by this gene is an axonal glycoprotein belonging to the immunoglobulin supergene family. The ectodomain, consisting of several immunoglobulin-like domains and fibronectin-like repeats (type III), is linked via a single transmembrane sequence to a conserved cytoplasmic domain. This cell adhesion molecule plays an important role in nervous system development, including neuronal migration and differentiation. Mutations in the gene cause X-linked neurological syndromes known as CRASH (corpus callosum hypoplasia, retardation, aphasia, spastic paraplegia and hydrocephalus). Alternative splicing of this gene results in multiple transcript variants, some of which include an alternate exon that is considered to be specific to neurons.

Protein Function: Neural cell adhesion molecule involved in the dynamics of cell adhesion and in the generation of transmembrane signals at tyrosine kinase receptors. During brain development, critical in multiple processes, including neuronal migration, axonal growth and fasciculation, and synaptogenesis. In the mature brain, plays a role in the dynamics of neuronal structure and function, including synaptic plasticity.

NCBI AND UNIPROT INFORMATION

NCBI GI #	4557707
NCBI GeneID	3897
NCBI Accession #	NP_000416.1
NCBI GenBank Nucleotide #	NM_000425.4
UniProt Accession #	P32004; Q8TA87; A0AV65; A4ZYW4; B2RMU7; G3XAF4
Molecular Weight	138,908 Da
NCBI Official Full Name	neural cell adhesion molecule L1 isoform 1
NCBI Official Synonym Full Names	L1 cell adhesion molecule
NCBI Official Symbol	L1CAM
NCBI Official Synonym Symbols	S10; HSAS; MASA; MIC5; SPG1; CAML1; CD171; HSAS1; N-CAML1; NCAM-L1; N-CAM-L1
NCBI Protein Information	neural cell adhesion molecule L1
UniProt Protein Name	Neural cell adhesion molecule L1
UniProt Gene Name	L1CAM

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/1cam-antibody-immunohistochemistry-elisa/19138>