

Rabbit CD105 Polyclonal Antibody | anti-CD105 antibody

Anti-CD105 Picoband Antibody


CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA19165	Rabbit	Mouse, Rat No cross reactivity with other proteins.	ELISA, Immunohistochemistry, Western Blot

PRODUCT SYNONYMS

CD105; Polyclonal Antibody; Anti-CD105 Picoband Antibody; Endoglin; Cell surface MJ7/18 antigen; Eng; Edg; anti-CD105 antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Mouse, Rat No cross reactivity with other proteins.
Clonality	Polyclonal
Purity/Purification	Immunogen affinity purified
Form/Format	Lyophilized
Concentration	Add 0.2ml of distilled water will yield a concentration of 500ug/ml. (varies by lot)
Sequence Length	653
Applicable Applications for anti-CD105 antibody	ELISA, IHC (Immunohistochemistry), WB (Western Blot)
Application Notes	WB: 0.1-0.5μg/ml Immunohistochemistry (Paraffin-embedded Section): 0.5-1μg/ml Direct ELISA: 0.1-0.5μg/ml
Immunogen	E Coli-derived mouse CD105 recombinant protein (Position: E27-L226).
Subcellular Localization	Membrane; Single-pass type I membrane protein.
Reconstitution	Add 0.2ml of distilled water will yield a concentration of 500ug/ml.
Contents	Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na2HPO4, 0.05mg NaN3.
Preparation and Storage	Store at -20 degree C for one year. After reconstitution, at 4 degree C for one month. It can also be aliquotted and stored frozen at -20 degree C for a longer time. Avoid repeated freezing and thawing.




IHC (Immunohistochemistry)

Figure 7. IHC analysis of CD105 using anti-CD105 antibody (AAA19165).CD105 was detected in paraffin-embedded section of rat spleen tissue. Heat mediated antigen retrieval was performed in citrate buffer (pH6, epitope retrieval solution) for 20 mins. The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1ug/ml rabbit anti-CD105 Antibody (AAA19165) overnight at 4 degree C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37 degree C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC) with DAB as the chromogen.




IHC (Immunohistochemistry)

Figure 6. IHC analysis of CD105 using anti-CD105 antibody (AAA19165).CD105 was detected in paraffin-embedded section of rat liver tissue. Heat mediated antigen retrieval was performed in citrate buffer (pH6, epitope retrieval solution) for 20 mins. The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1ug/ml rabbit anti-CD105 Antibody (AAA19165) overnight at 4 degree C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37 degree C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC) with DAB as the chromogen.




IHC (Immunohistochemistry)

Figure 5. IHC analysis of CD105 using anti-CD105 antibody (AAA19165).CD105 was detected in paraffin-embedded section of rat kidney tissue. Heat mediated antigen retrieval was performed in citrate buffer (pH6, epitope retrieval solution) for 20 mins. The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1ug/ml rabbit anti-CD105 Antibody (AAA19165) overnight at 4 degree C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37 degree C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC) with DAB as the chromogen.




IHC (Immunohistochemistry)

Figure 4. IHC analysis of CD105 using anti-CD105 antibody (AAA19165).CD105 was detected in paraffin-embedded section of mouse spleen tissue. Heat mediated antigen retrieval was performed in citrate buffer (pH6, epitope retrieval solution) for 20 mins. The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1ug/ml rabbit anti-CD105 Antibody (AAA19165) overnight at 4 degree C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37 degree C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC) with DAB as the chromogen.



IHC (Immunohistochemistry)

Figure 3. IHC analysis of CD105 using anti-CD105 antibody (AAA19165).CD105 was detected in paraffin-embedded section of mouse liver tissue. Heat mediated antigen retrieval was performed in citrate buffer (pH6, epitope retrieval solution) for 20 mins. The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1ug/ml rabbit anti-CD105 Antibody (AAA19165) overnight at 4 degree C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37 degree C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC) with DAB as the chromogen.



IHC (Immunohistochemistry)

Figure 2. IHC analysis of CD105 using anti-CD105 antibody (AAA19165).CD105 was detected in paraffin-embedded section of mouse kidney tissue. Heat mediated antigen retrieval was performed in citrate buffer (pH6, epitope retrieval solution) for 20 mins. The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1ug/ml rabbit anti-CD105 Antibody (AAA19165) overnight at 4 degree C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37 degree C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC) with DAB as the chromogen.

ADDITIONAL INFORMATION

Related Product Information for anti-CD105 antibody	Description: Endoglin (Osler-Rendu-Weber syndrome 1), CD105, is a type I membrane glycoprotein located on cell surfaces and is a part of the TGF beta receptor complex. Its gene is mapped to human chromosome 8. The protein consists of a homodimer of 180 kDA with disulfide links. It has been found on endothelial cells, activated macrophages, fibroblasts and smooth muscle cells. Endoglin has a role in the development of the cardiovascular system and in vascular remodeling and has been found to be elevated in pregnant women who subsequently develop preeclampsia. Protein Function: Major glycoprotein of vascular endothelium. Involved in the regulation of angiogenesis. May play a critical role in the binding of endothelial cells to integrins and/or other RGD receptors. Acts as TGF-beta coreceptor and is involved in the TGF- beta/BMP signaling cascade. Required for GDF2/BMP9 signaling through SMAD1 in endothelial cells.
Product Categories/Family for anti-CD105 antibody	Neuroscience; Neurology Process; Neurogenesis; Epigenetics and Nuclear Signaling; Transcription; Other factors

NCBI AND UNIPROT INFORMATION

NCBI GI #	226437647
NCBI GeneID	13805
NCBI Accession #	NP_031958.2
NCBI GenBank Nucleotide #	NM_007932.2
UniProt Accession #	Q63961; Q61520; Q8K100
Molecular Weight	70,021 Da
NCBI Official Full Name	endoglin isoform 1
NCBI Official Synonym Full Names	endoglin
NCBI Official Symbol	Eng
NCBI Official Synonym Symbols	Endo; CD105; AI528660; AI662476; S-endoglin
NCBI Protein Information	endoglin
UniProt Protein Name	Endoglin
UniProt Gene Name	Eng
UniProt Synonym Gene Names	Edg

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/cd105-antibody-immunohistochemistry-western-blot-elisa/19165>