

Rabbit Annexin VI Polyclonal Antibody | anti-ANXA6 antibody

Anti-Annexin VI Picoband Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA19170	Rabbit	Human, Mouse, Rat No cross reactivity with other proteins.	ELISA, Immunohistochemistry, Western Blot

PRODUCT SYNONYMS

Annexin VI; Polyclonal Antibody; Anti-Annexin VI Picoband Antibody; Annexin A6; 67 kDa calelectrin; Annexin-6; Calphobindin-II; CPB-II; Chromobindin-20; Lipocortin VI; Protein III; p68; p70; ANXA6; ANX6; anti-ANXA6 antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat No cross reactivity with other proteins.
Clonality	Polyclonal
Form/Format	Lyophilized
Sequence Length	641
Applicable Applications for anti-ANXA6 antibody	ELISA, IHC (Immunohistochemistry), WB (Western Blot)
Application Notes	WB: 0.1-0.5µg/ml Immunohistochemistry (Paraffin-embedded Section): 0.5-1µg/ml Direct ELISA: 0.1-0.5µg/ml
Immunogen	E Coli-derived human Annexin VI recombinant protein (Position: N395-L665).
Subcellular Localization	Cytoplasm. Melanosome
Reconstitution	Add 0.2ml of distilled water will yield a concentration of 500µg/ml.
Contents	Each vial contains 4mg Trehalose, 0.9mg NaCl, 0.2mg Na2HPO4, 0.05mg NaN3.
Preparation and Storage	Store at -20 degree C for one year. After reconstitution, at 4 degree C for one month. It can also be aliquotted and stored frozen at -20 degree C for a longer time. Avoid repeated freezing and thawing.

IMAGES



IHC (Immunohistochemistry)

Figure 6. IHC analysis of Annexin VI using anti-Annexin VI antibody (AAA19170). Annexin VI was detected in paraffin-embedded section of rat spleen tissue. Heat mediated antigen retrieval was performed in citrate buffer (pH6, epitope retrieval solution) for 20 mins. The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2µg/ml rabbit anti-Annexin VI Antibody (AAA19170) overnight at 4 degree C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37 degree C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC) with DAB as the chromogen.



IHC (Immunohistochemistry)

Figure 5. IHC analysis of Annexin VI using anti-Annexin VI antibody (AAA19170). Annexin VI was detected in paraffin-embedded section of mouse spleen tissue. Heat mediated antigen retrieval was performed in citrate buffer (pH6, epitope retrieval solution) for 20 mins. The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2µg/ml rabbit anti-Annexin VI Antibody (AAA19170) overnight at 4 degree C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37 degree C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC) with DAB as the chromogen.




IHC (Immunohistochemistry)

Figure 4. IHC analysis of Annexin VI using anti-Annexin VI antibody (AAA19170).Annexin VI was detected in paraffin-embedded section of human lung cancer tissue. Heat mediated antigen retrieval was performed in citrate buffer (pH6, epitope retrieval solution) for 20 mins. The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2ug/ml rabbit anti-Annexin VI Antibody (AAA19170) overnight at 4 degree C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37 degree C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC) with DAB as the chromogen.




IHC (Immunohistochemistry)

Figure 3. IHC analysis of Annexin VI using anti-Annexin VI antibody (AAA19170).Annexin VI was detected in paraffin-embedded section of human intestinal cancer tissue. Heat mediated antigen retrieval was performed in citrate buffer (pH6, epitope retrieval solution) for 20 mins. The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2ug/ml rabbit anti-Annexin VI Antibody (AAA19170) overnight at 4 degree C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37 degree C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC) with DAB as the chromogen.



WB (Western Blot)

Figure 2. Western blot analysis of Annexin VI using anti-Annexin VI antibody (AAA19170).Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 50ug of sample under reducing conditions.Lane 1: human Hela whole cell lysates,Lane 2: human placenta tissue lysates,Lane 3: human MCF-7 whole cell lysates,Lane 4: human U-87MG whole cell lysates,Lane 5: human HepG2 whole cell lysates,Lane 6: human SMMC-7721 whole cell lysates,Lane 7: human 293T whole cell lysates.After Electrophoresis, proteins were transferred to a Nitrocellulose membrane at 150mA for 50-90 minutes. Blocked the membrane with 5% Non-fat Milk/ TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-Annexin VI antigen affinity purified polyclonal antibody at 0.5ug/mL overnight at 4 degree C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:10000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit with Tanon 5200 system. A specific band was detected for Annexin VI at approximately 76KD. The expected band size for Annexin VI is at 76KD.



WB (Western Blot)

Figure 1. Western blot analysis of Annexin VI using anti-Annexin VI antibody (AAA19170).Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 50ug of sample under reducing conditions.Lane 1: rat brain tissue lysates,Lane 2: rat thymus tissue lysates,Lane 3: rat testis tissue lysates,Lane 4: rat pancreas tissue lysates,Lane 5: mouse brain tissue lysates,Lane 6: mouse thymus tissue lysates,Lane 7: mouse testis tissue lysates,Lane 8: mouse pancreas tissue lysates,Lane 9: mouse HEPA1-6 whole cell lysates.After Electrophoresis, proteins were transferred to a Nitrocellulose membrane at 150mA for 50-90 minutes. Blocked the membrane with 5% Non-fat Milk/ TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-Annexin VI antigen affinity purified polyclonal antibody at 0.5ug/mL overnight at 4 degree C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:10000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit with Tanon 5200 system. A specific band was detected for Annexin VI at approximately 76KD. The expected band size for Annexin VI is at 76KD.

ADDITIONAL INFORMATION

Related Product Information for anti-ANXA6 antibody

Description: Annexin A6 (ANXA6) is a member of a family of proteins that bind membrane or cytoskeleton in a Ca(2+)-dependent manner. These proteins are characterized by homologous amino acid sequences that are present in multiple copies in each protein. ANXA6 gene is assigned to 5q32-q34 by use of a cDNA clone to probe genomic DNA from rodent-human somatic cell hybrids and for in situ hybridization. The ANX6 gene is approximately 60 kb long and contains 26 exons. The genomic sequence at the 3-prime end does not contain a canonical polyadenylation signal. Ca(2+)-dependent binding between CRHSP28 and ANXA6 is required for acinar cell membrane trafficking events and digestive enzyme secretion.
Protein Function: May associate with CD21. May regulate the release of Ca(2+) from intracellular stores.

NCBI AND UNIPROT INFORMATION

NCBI GI #	71773329
NCBI GeneID	309
NCBI Accession #	NP_001146.2
NCBI GenBank Nucleotide #	NM_001155.4
UniProt Accession #	P08133 ; Q6ZT79 ; B7Z8A7 ; D3DQH4 ; E9PGK1
Molecular Weight	72,423 Da
NCBI Official Full Name	annexin A6 isoform 1
NCBI Official Synonym Full Names	annexin A6
NCBI Official Symbol	ANXA6

NCBI Official Synonym Symbols	ANX6; CBP68
NCBI Protein Information	annexin A6
UniProt Protein Name	Annexin A6
UniProt Gene Name	ANXA6
UniProt Synonym Gene Names	ANX6; CPB-II

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/annexin-vi-antibody-immunohistochemistry-western-blot-elisa/19170>