

Rabbit MEK2/MAP2K2 Polyclonal Antibody | anti-MAP2K2 antibody

Anti-MEK2/MAP2K2 Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA19230	Rabbit	Human, Mouse, Rat	Western Blot, Immunohistochemistry, Immunocytochemistry, Immunofluorescence, Flow Cytometry, Functional Assay, ELISA

PRODUCT SYNONYMS


MEK2/MAP2K2; Polyclonal Antibody; Anti-MEK2/MAP2K2 Antibody; MAP2K2; MEK2; MKK2; PRKMK2; Dual specificity mitogen-activated protein kinase kinase 2; MAP kinase kinase 2; MAPKK 2; EC 2. 7. 12. 2; ERK activator kinase 2; MAPK/ERK kinase 2; MEK 2; mitogen-activated protein kinase kinase 2; anti-MAP2K2 antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Isotype	Rabbit IgG
Specificity	Rabbit IgG polyclonal antibody for MEK2/MAP2K2 detection.
Purity/Purification	Immunogen affinity purified.
Form/Format	Lyophilized. Each vial contains 4mg Trehalose, 0.9mg NaCl, 0.2mg Na2HPO4, 0.01mg NaN3.
Applicable Applications for anti-MAP2K2 antibody	WB (Western Blot), IHC (Immunohistochemistry), ICC (Immunocytochemistry), IF (Immunofluorescence), FCM/FACS (Flow Cytometry), ELISA (Direct ELISA)
Application Notes	WB: 0.25-0.5ug/ml Human, Mouse, Rat IHC-P: 2-5ug/ml Human, Mouse, Rat ICC/IF: 5ug/ml Human FC/FACS/FCM: 1-3ug/1x106 cells Human Direct ELISA: 0.1-0.5ug/ml Human
Immunogen	E Coli-derived human MEK2/MAP2K2 recombinant protein (Position: M1-V400).
Reconstitution	Add 0.2ml of distilled water will yield a concentration of 500ug/ml.
Recommended Detection Systems	Recommended Detection Systems
Preparation and Storage	Store at -20 degree C for one year. After reconstitution, at 4 degree C for one month. It can also be aliquotted and stored frozen at -20 degree C for a longer time. Avoid repeated freezing and thawing.




IHC (Immunohistochemistry)
Figure 8. IHC analysis of MEK2/MAP2K2 using anti-MEK2/MAP2K2 antibody (AAA19230).MEK2/MAP2K2 was detected in paraffin-embedded section of mouse brain tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH8. 0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2μg/ml rabbit anti-MEK2/MAP2K2 Antibody (AAA19230) overnight at 4 degree C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37 degree C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC) (Catalog # with DAB as the chromogen.



FCM (Flow Cytometry)
Figure 7. Flow Cytometry analysis of SiHa cells using anti-MEK2/MAP2K2 antibody (AAA19230).Overlay histogram showing PC-3 cells stained with AAA19230 (Blue line). The cells were blocked with 10% normal goat serum. And then incubated with rabbit anti-MEK2/MAP2K2 Antibody (AAA19230, 1μg/1x10⁶ cells) for 30 min at 20 degree C. DyLight®488 conjugated goat anti-rabbit IgG (5-10μg/1x10⁶ cells) was used as secondary antibody for 30 minutes at 20 degree C. Isotype control antibody (Green line) was rabbit IgG (1μg/1x10⁶) used under the same conditions. Unlabelled sample (Red line) was also used as a control.



IF (Immunofluorescence)
Figure 6. IF analysis of MEK2/MAP2K2 using anti- MEK2/MAP2K2 antibody (AAA19230).MEK2/MAP2K2 was detected in immunocytochemical section of K562 cells. Enzyme antigen retrieval was performed using IHC enzyme antigen retrieval reagent for 15 mins. The cells were blocked with 10% goat serum. And then incubated with 5μg/mL rabbit anti- MEK2/MAP2K2 Antibody (AAA19230) overnight at 4 degree C. DyLight®594 Conjugated Goat Anti-Rabbit IgG (BA1142) was used as secondary antibody at 1:100 dilution and incubated for 30 minutes at 37 degree C. The section was counterstained with DAPI. Visualize using a fluorescence microscope and filter sets appropriate for the label used.



IF (Immunofluorescence)
Figure 5. IF analysis of MEK2/MAP2K2 using anti- MEK2/MAP2K2 antibody (AAA19230).MEK2/MAP2K2 was detected in immunocytochemical section of U2OS cells. Enzyme antigen retrieval was performed using IHC enzyme antigen retrieval reagent for 15 mins. The cells were blocked with 10% goat serum. And then incubated with 5μg/mL rabbit anti- MEK2/MAP2K2 Antibody (AAA19230) overnight at 4 degree C. DyLight®488 Conjugated Goat Anti-Rabbit IgG was used as secondary antibody at 1:100 dilution and incubated for 30 minutes at 37 degree C. The section was counterstained with DAPI. Visualize using a fluorescence microscope and filter sets appropriate for the label used.



IHC (Immunohistochemistry)
Figure 4. IHC analysis of MEK2/MAP2K2 using anti-MEK2/MAP2K2 antibody (AAA19230).MEK2/MAP2K2 was detected in paraffin-embedded section of rat brain tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH8. 0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2μg/ml rabbit anti-MEK2/MAP2K2 Antibody (AAA19230) overnight at 4 degree C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37 degree C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC) (Catalog # with DAB as the chromogen.



IHC (Immunohistochemistry)
Figure 3. IHC analysis of MEK2/MAP2K2 using anti-MEK2/MAP2K2 antibody (AAA19230).MEK2/MAP2K2 was detected in paraffin-embedded section of rat brain tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH8. 0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2μg/ml rabbit anti-MEK2/MAP2K2 Antibody (AAA19230) overnight at 4 degree C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37 degree C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC) (Catalog # with DAB as the chromogen.

ADDITIONAL INFORMATION

Related Product Information for anti-MAP2K2 antibody	Dual specificity mitogen-activated protein kinase kinase 2(MAP2K2), also called PRKMK2 or MEK2, is an enzyme that in humans is encoded by the MAP2K2 gene. The protein encoded by this gene is a dual specificity protein kinase that belongs to the MAP kinase kinase family. MAP2K2 is mapped to 19p13. 3. This kinase is known to play a critical role in mitogen growth factor signal transduction, and the inhibition or degradation of this kinase is found to be involved in the pathogenesis of Yersinia and anthrax. Recombinant MEK2 and MEK1 both could activate human ERK1 in vitro, and they further characterized biochemically the 2 MAP2Ks. MAP2K2 has been shown to interact with MAPK3 and ARAF.
References	1. "Entrez Gene: MAP2K2 mitogen-activated protein kinase kinase 2" 2. Zheng, C. F., Guan, K. L. Cloning and characterization of two distinct human extracellular signal-regulated kinase activator kinases, MEK1 and MEK2. J. Biol. Chem. 268: 11435-11439, 1993.

NCBI AND UNIPROT INFORMATION

NCBI GI #	547915
NCBI GeneID	5605
UniProt Accession #	P36507

Molecular Weight	400
NCBI Official Full Name	Dual specificity mitogen-activated protein kinase kinase 2
NCBI Official Synonym Full Names	mitogen-activated protein kinase kinase 2
NCBI Official Symbol	MAP2K2
NCBI Official Synonym Symbols	CFC4; MEK2; MKK2; MAPKK2; PRKMK2
NCBI Protein Information	dual specificity mitogen-activated protein kinase kinase 2; MAPK/ERK kinase 2; MAP kinase kinase 2; ERK activator kinase 2; mitogen-activated protein kinase kinase 2, p45
UniProt Protein Name	Dual specificity mitogen-activated protein kinase kinase 2
UniProt Gene Name	MAP2K2
UniProt Synonym Gene Names	MEK2; MKK2; PRKMK2; MAP kinase kinase 2; MAPKK 2; MEK 2
UniProt Entry Name	MP2K2_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/mek2-map2k2-antibody-western-blot-immunohistochemistry-immunocytochemistry-immunofluorescence-flow-cytometry-direct-elisa-elisa/19230>