

Rabbit GNG2 Polyclonal Antibody | anti-GNG2 antibody

Anti-GNG2 Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA19312	Rabbit	Human, Mouse, Rat	Western Blot, Immunohistochemistry, Flow Cytometry, Functional Assay, ELISA


PRODUCT SYNONYMS

GNG2; Polyclonal Antibody; Anti-GNG2 Antibody; Gng2; Guanine nucleotide-binding protein G(I)/G(S)/G(O subunit gamma-2; G gamma-I; G protein subunit gamma 2; anti-GNG2 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Isotype	Rabbit IgG
Specificity	Rabbit IgG polyclonal antibody for GNG2 detection.
Purity/Purification	Immunogen affinity purified.
Form/Format	Lyophilized. Each vial contains 4mg Trehalose, 0.9mg NaCl, 0.2mg Na2HPO4, 0.01mg NaN3.
Applicable Applications for anti-GNG2 antibody	WB (Western Blot), IHC (Immunohistochemistry), FCM/FACS (Flow Cytometry), ELISA (Direct ELISA)
Application Notes	WB: 0.1-0.25ug/ml Human, Mouse, Rat IHC-P: 2-5ug/ml Human FC/FACS/FCM: 1-3ug/1x106 cells Human, Mouse Direct ELISA: 0.1-0.5ug/ml Human
Immunogen	E Coli-derived human GNG2 recombinant protein (Position: A2-D48).
Reconstitution	Add 0.2ml of distilled water will yield a concentration of 500ug/ml.
Recommended Detection Systems	Recommended Detection Systems
Preparation and Storage	Store at -20 degree C for one year. After reconstitution, at 4 degree C for one month. It can also be aliquotted and stored frozen at -20 degree C for a longer time. Avoid repeated freezing and thawing.

IMAGES



FCM (Flow Cytometry)

Figure 10. Flow Cytometry analysis of U87 cells using anti-GNG2 antibody (AAA19312). Overlay histogram showing U87 cells stained with AAA19312 (Blue line). The cells were blocked with 10% normal goat serum. And then incubated with rabbit anti-GNG2 Antibody (AAA19312, 1μg/1x106 cells) for 30 min at 20 degree C. DyLight®488 conjugated goat anti-rabbit IgG (5-10μg/1x106 cells) was used as secondary antibody for 30 minutes at 20 degree C. Isotype control antibody (Green line) was rabbit IgG (1μg/1x106) used under the same conditions. Unlabelled sample (Red line) was also used as a control.




FCM (Flow Cytometry)

Figure 9. Flow Cytometry analysis of HEPA1-6 cells using anti-GNG2 antibody (AAA19312). Overlay histogram showing HEPA1-6 cells stained with AAA19312 (Blue line). The cells were blocked with 10% normal goat serum. And then incubated with rabbit anti-GNG2 Antibody (AAA19312, 1μg/1x106 cells) for 30 min at 20 degree C. DyLight®488 conjugated goat anti-rabbit IgG (5-10μg/1x106 cells) was used as secondary antibody for 30 minutes at 20 degree C. Isotype control antibody (Green line) was rabbit IgG (1μg/1x106) used under the same conditions. Unlabelled sample (Red line) was also used as a control.




IHC (Immunohistochemistry)

Figure 8. IHC analysis of GNG2 using anti-GNG2 antibody (AAA19312). GNG2 was detected in paraffin-embedded section of human renal carcinoma tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH8. 0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2µg/ml rabbit anti-GNG2 Antibody (AAA19312) overnight at 4 degree C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37 degree C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC) (Catalog # with DAB as the chromogen).




IHC (Immunohistochemistry)

Figure 7. IHC analysis of GNG2 using anti-GNG2 antibody (AAA19312). GNG2 was detected in paraffin-embedded section of human placenta tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH8. 0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2µg/ml rabbit anti-GNG2 Antibody (AAA19312) overnight at 4 degree C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37 degree C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC) (Catalog # with DAB as the chromogen).



IHC (Immunohistochemistry)

Figure 6. IHC analysis of GNG2 using anti-GNG2 antibody (AAA19312). GNG2 was detected in paraffin-embedded section of human ovarian cancer tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH8. 0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2µg/ml rabbit anti-GNG2 Antibody (AAA19312) overnight at 4 degree C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37 degree C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC) (Catalog # with DAB as the chromogen).



IHC (Immunohistochemistry)

Figure 5. IHC analysis of GNG2 using anti-GNG2 antibody (AAA19312). GNG2 was detected in paraffin-embedded section of human lung cancer tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH8. 0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2µg/ml rabbit anti-GNG2 Antibody (AAA19312) overnight at 4 degree C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37 degree C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC) (Catalog # with DAB as the chromogen).

ADDITIONAL INFORMATION

Related Product Information for anti-GNG2 antibody	Guanine nucleotide-binding protein G(I)/G(S)/G(O) subunit gamma-2 is a protein that in humans is encoded by the GNG2 gene. This gene encodes one of the gamma subunits of a guanine nucleotide-binding protein. Such proteins are involved in signaling mechanisms across membranes. Various subunits forms heterodimers which then interact with the different signal molecules.
References	<ol style="list-style-type: none">Downes, G. B., Gilbert, D. J., Copeland, N. G., Gautam, N., Jenkins, N. A. Chromosomal mapping of five mouse G protein gamma subunits. <i>Genomics</i> 57: 173-176, 1999.Modarressi, M. H., Taylor, K. E., Wolfe, J. Cloning, characterization, and mapping of the gene encoding the human G protein gamma-2 subunit. <i>Biochem. Biophys. Res. Commun.</i> 272: 610-615, 2000.Wolfe, J. T., Wang, H., Howard, J., Garrison, J. C., Barrett, P. Q. T-type calcium channel regulation by specific G-protein beta-gamma subunits. <i>Nature</i> 424: 209-213, 2003.

NCBI AND UNIPROT INFORMATION

NCBI GI #	344313196
NCBI GeneID	54331
NCBI Accession #	NP_001230702.1
NCBI GenBank Nucleotide #	NM_001243773.1
UniProt Accession #	P59768 ; Q5JPE2 ; Q6P9A9
Molecular Weight	7,850 Da
NCBI Official Full Name	guanine nucleotide-binding protein G(I)/G(S)/G(O) subunit gamma-2
NCBI Official Synonym Full Names	guanine nucleotide binding protein (G protein), gamma 2
NCBI Official Symbol	GNG2
NCBI Protein Information	guanine nucleotide-binding protein G(I)/G(S)/G(O) subunit gamma-2; guanine nucleotide-binding protein G(I)/G(S)/G(O) subunit gamma-2; g gamma-I; guanine nucleotide binding protein gamma 2; guanine nucleotide-binding protein G(I)/G(O) gamma-2 subunit

UniProt Protein Name	Guanine nucleotide-binding protein G(I)/G(S)/G(O) subunit gamma-2
UniProt Gene Name	GNG2
UniProt Entry Name	GBG2_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/gng2-antibody-western-blot-immunohistochemistry-flow-cytometry-direct-elisa-elisa/19312>