

Rabbit anti-Human IMPDH2 Polyclonal Antibody | anti-IMPDH2 antibody

Anti-IMPDH2 Antibody Picoband

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA19475	Rabbit	Human	Flow Cytometry, Functional Assay, Immunofluorescence, Immunohistochemistry, Immunocytochemistry, Western Blot

PRODUCT SYNONYMS

IMPDH2; Polyclonal Antibody; Anti-IMPDH2 Antibody Picoband; anti-IMPDH2 antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	Rabbit IgG
Purity/Purification	Immunogen affinity purified.
Form/Format	Lyophilized Each vial contains 4 mg Trehalose, 0.9 mg NaCl, 0.2 mg Na ₂ HPO ₄ .
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml. (varies by lot)
Applicable Applications for anti-IMPDH2 antibody	FCM/FACS (Flow Cytometry), IF (Immunofluorescence), IHC (Immunohistochemistry), ICC (Immunocytochemistry), WB (Western Blot)
Application Notes	WB: 0.25-0.5 ug/ml/ml, Human IHC-P: 2-5 ug/ml/ml, Human ICC/IF: 5 ug/ml, Human FC/FACS: 1-3 ug/ml/1x10 ⁶ cells, Human Tested Species: In-house tested species with positive results. Enhanced Chemiluminescent Kit with anti-Rabbit IgG for Western blot, and HRP Conjugated anti-Rabbit IgG Super Vision Assay Kit for IHC(P) and ICC.
Reconstitution	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Immunogen	A synthetic peptide corresponding to a sequence of human IMPDH2 (VVLSPKDRVRDVFEAKARH).
Preparation and Storage	Store at -20 degree C for one year from date of receipt. After reconstitution, at 4 degree C for one month. It can also be aliquotted and stored frozen at -20 degree C for six months. Avoid repeated freezing and thawing.



FCM (Flow Cytometry)

Figure 7. Flow Cytometry analysis of U2OS cells using anti-IMPDPH2 antibody (AAA19475).Overlay histogram showing U2OS cells stained with AAA19475 (Blue line). The cells were blocked with 10% normal goat serum. And then incubated with rabbit anti-IMPDPH2 Antibody (AAA19475, 1 ug/1x10⁶ cells) for 30 min at 20 degree C. DyLight488 conjugated goat anti-rabbit IgG was used as secondary antibody for 30 minutes at 20 degree C. Isotype control antibody (Green line) was rabbit IgG (1 ug/1x10⁶) used under the same conditions. Unlabelled sample (Red line) was also used as a control.



IF (Immunofluorescence)

Figure 6. IF analysis of IMPDPH2 using anti-IMPDPH2 antibody (AAA19475).IMPDPH2 was detected in an immunocytochemical section of T47D cells. Enzyme antigen retrieval was performed using IHC enzyme antigen retrieval reagent (AR0022) for 15 mins. The cells were blocked with 10% goat serum. And then incubated with 5 ug/mL rabbit anti-IMPDPH2 Antibody (AAA19475) overnight at 4 degree C. DyLight594 Conjugated Goat Anti-Rabbit IgG (BA1142) was used as secondary antibody at 1:100 dilution and incubated for 30 minutes at 37 degree C. The section was counterstained with DAPI. Visualize using a fluorescence microscope and filter sets appropriate for the label used.



IHC (Immunohistochemistry)

Figure 5. IHC analysis of IMPDPH2 using anti-IMPDPH2 antibody (AAA19475).IMPDPH2 was detected in a paraffin-embedded section of human ovarian cancer tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 ug/ml rabbit anti-IMPDPH2 Antibody (AAA19475) overnight at 4 degree C. Peroxidase Conjugated Goat Anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37 degree C. The tissue section was developed using HRP Conjugated Rabbit IgG Super Vision Assay Kit with DAB as the chromogen.



IHC (Immunohistochemistry)

Figure 4. IHC analysis of IMPDPH2 using anti-IMPDPH2 antibody (AAA19475).IMPDPH2 was detected in a paraffin-embedded section of human cervical cancer tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 ug/ml rabbit anti-IMPDPH2 Antibody (AAA19475) overnight at 4 degree C. Peroxidase Conjugated Goat Anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37 degree C. The tissue section was developed using HRP Conjugated Rabbit IgG Super Vision Assay Kit with DAB as the chromogen.



IHC (Immunohistochemistry)

Figure 3. IHC analysis of IMPDPH2 using anti-IMPDPH2 antibody (AAA19475).IMPDPH2 was detected in a paraffin-embedded section of human breast cancer tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 ug/ml rabbit anti-IMPDPH2 Antibody (AAA19475) overnight at 4 degree C. Peroxidase Conjugated Goat Anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37 degree C. The tissue section was developed using HRP Conjugated Rabbit IgG Super Vision Assay Kit with DAB as the chromogen.



IHC (Immunohistochemistry)

Figure 2. IHC analysis of IMPDPH2 using anti-IMPDPH2 antibody (AAA19475).IMPDPH2 was detected in a paraffin-embedded section of human esophageal squamous carcinoma tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 ug/ml rabbit anti-IMPDPH2 Antibody (AAA19475) overnight at 4 degree C. Peroxidase Conjugated Goat Anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37 degree C. The tissue section was developed using HRP Conjugated Rabbit IgG Super Vision Assay Kit with DAB as the chromogen.

ADDITIONAL INFORMATION

Related Product Information for anti-IMPDPH2 antibody	Inosine-5'-monophosphate dehydrogenase 2, also known as IMP dehydrogenase 2, is an enzyme that in humans is encoded by the IMPDPH2 gene. This gene encodes the rate-limiting enzyme in the de novo guanine nucleotide biosynthesis. It is thus involved in maintaining cellular guanine deoxy- and ribonucleotide pools needed for DNA and RNA synthesis. The encoded protein catalyzes the NAD-dependent oxidation of inosine-5'-monophosphate into xanthine-5'-monophosphate, which is then converted into guanosine-5'-monophosphate. This gene is up-regulated in some neoplasms, suggesting it may play a role in malignant transformation.
Product Categories/Family for anti-IMPDPH2 antibody	Primary Antibodies, Rabbit Polyclonal Antibodies

NCBI AND UNIPROT INFORMATION

NCBI GI #	66933016
NCBI GeneID	3615
NCBI Accession #	NP_000875.2
NCBI GenBank Nucleotide #	NM_000884.2

UniProt Accession #	P12268; Q6LEF3
Molecular Weight	55,805 Da
NCBI Official Full Name	inosine-5'-monophosphate dehydrogenase 2
NCBI Official Synonym Full Names	IMP (inosine 5'-monophosphate) dehydrogenase 2
NCBI Official Symbol	IMPDH2
NCBI Official Synonym Symbols	IMPD2; IMPDH-II
NCBI Protein Information	inosine-5'-monophosphate dehydrogenase 2; IMP (inosine monophosphate) dehydrogenase 2; IMP oxireductase 2; IMPD 2; IMPDH 2; IMPDH2; inosine 5' phosphate dehydrogenase 2; inosine monophosphate dehydrogenase type II
UniProt Protein Name	Inosine-5'-monophosphate dehydrogenase 2
UniProt Gene Name	IMPDH2
UniProt Entry Name	IMDH2_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/impdh2-antibody-flow-cytometry-immunofluorescence-immunohistochemistry-immunocytochemistry-western-blot-elisa/19475>