

Rabbit EIF5 Polyclonal Antibody | anti-EIF5 antibody

Anti-EIF5 Antibody Picoband


CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA19525	Rabbit	Human, Mouse, Rat	Western Blot, Immunohistochemistry, Flow Cytometry, Functional Assay, ELISA

PRODUCT SYNONYMS

EIF5; Polyclonal Antibody; Anti-EIF5 Antibody Picoband; anti-EIF5 antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Isotype	Rabbit IgG
Purity/Purification	Immunogen affinity purified.
Form/Format	Lyophilized Each vial contains 4 mg Trehalose, 0.9 mg NaCl, 0.2 mg Na ₂ HPO ₄ .
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml. (varies by lot)
Applicable Applications for anti-EIF5 antibody	WB (Western Blot), IHC (Immunohistochemistry), FCM/FACS (Flow Cytometry), ELISA
Application Notes	WB: 0.1-0.25 ug/ml, Human IHC-P: 2-5 ug/ml, Human, Mouse, Rat FC/FACS: 1-3 ug/1x10 ⁶ cells, Human Direct ELISA: 0.1-0.5 ug/ml, Human Tested Species: In-house tested species with positive results. Enhanced Chemiluminescent Kit with anti-Rabbit IgG for Western blot, and HRP Conjugated anti-Rabbit IgG Super Vision Assay Kit for IHC(P).
Reconstitution	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Immunogen	E Coli-derived human EIF5 recombinant protein (Position: D11-D423).
Preparation and Storage	Store at -20 degree C for one year from date of receipt. After reconstitution, at 4 degree C for one month. It can also be aliquotted and stored frozen at -20 degree C for six months. Avoid repeated freezing and thawing.



FCM (Flow Cytometry)

Figure 12. Flow Cytometry analysis of K562 cells using anti-EIF5 antibody (AAA19525).Overlay histogram showing K562 cells stained with AAA19525 (Blue line). The cells were blocked with 10% normal goat serum. And then incubated with rabbit anti-EIF5 Antibody (AAA19525, 1 ug/1x10⁶ cells) for 30 min at 20 degree C. DyLight488 conjugated goat anti-rabbit IgG (BA1127, 5-10 ug/1x10⁶ cells) was used as secondary antibody for 30 minutes at 20 degree C. Isotype control antibody (Green line) was rabbit IgG (1 ug/1x10⁶) used under the same conditions. Unlabelled sample (Red line) was also used as a control.




IHC (Immunohistochemistry)

Figure 10. IHC analysis of EIF5 using anti-EIF5 antibody (AAA19525).EIF5 was detected in a paraffin-embedded section of mouse brain tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 ug/ml rabbit anti-EIF5 Antibody (AAA19525) overnight at 4 degree C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37 degree C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC) with DAB as the chromogen.




IHC (Immunohistochemistry)

Figure 11. IHC analysis of EIF5 using anti-EIF5 antibody (AAA19525).EIF5 was detected in a paraffin-embedded section of rat brain tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 ug/ml rabbit anti-EIF5 Antibody (AAA19525) overnight at 4 degree C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37 degree C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC) with DAB as the chromogen.




IHC (Immunohistochemistry)

Figure 9. IHC analysis of EIF5 using anti-EIF5 antibody (AAA19525).EIF5 was detected in a paraffin-embedded section of mouse brain tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 ug/ml rabbit anti-EIF5 Antibody (AAA19525) overnight at 4 degree C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37 degree C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC) with DAB as the chromogen.



IHC (Immunohistochemistry)

Figure 8. IHC analysis of EIF5 using anti-EIF5 antibody (AAA19525). EIF5 was detected in a paraffin-embedded section of human renal cancer tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 ug/ml rabbit anti-EIF5 Antibody (AAA19525) overnight at 4 degree C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37 degree C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC) with DAB as the chromogen.



IHC (Immunohistochemistry)

Figure 7. IHC analysis of EIF5 using anti-EIF5 antibody (AAA19525). EIF5 was detected in a paraffin-embedded section of human placenta tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 ug/ml rabbit anti-EIF5 Antibody (AAA19525) overnight at 4 degree C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37 degree C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC) with DAB as the chromogen.

ADDITIONAL INFORMATION

Related Product Information for anti-EIF5 antibody	Eukaryotic translation initiation factor 5 is a protein that in humans is encoded by the EIF5 gene. Eukaryotic translation initiation factor-5 (EIF5) interacts with the 40S initiation complex to promote hydrolysis of bound GTP with concomitant joining of the 60S ribosomal subunit to the 40S initiation complex. The resulting functional 80S ribosomal initiation complex is then active in peptidyl transfer and chain elongations.
Product Categories/Family for anti-EIF5 antibody	Primary Antibodies , Rabbit Polyclonal Antibodies

NCBI AND UNIPROT INFORMATION

NCBI GI #	21361337
NCBI GeneID	1983
NCBI Accession #	NP_001960
NCBI GenBank Nucleotide #	NM_001969
UniProt Accession #	P55010 ; Q53XB3 ; Q9H5N2 ; Q9UG48
Molecular Weight	49,223 Da
NCBI Official Full Name	eukaryotic translation initiation factor 5
NCBI Official Synonym Full Names	eukaryotic translation initiation factor 5
NCBI Official Symbol	EIF5
NCBI Official Synonym Symbols	EIF-5; EIF-5A
NCBI Protein Information	eukaryotic translation initiation factor 5
UniProt Protein Name	Eukaryotic translation initiation factor 5
UniProt Gene Name	EIF5
UniProt Synonym Gene Names	eIF-5
UniProt Entry Name	IF5_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/eif5-antibody-western-blot-immunohistochemistry-flow-cytometry-elisa/19525>