

Rabbit MDH2 Polyclonal Antibody | anti-MDH2 antibody

Anti-MDH2 Antibody Picoband


CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA19527	Rabbit	Human, Mouse, Rat	Western Blot, Immunohistochemistry, Immunocytochemistry, Immunofluorescence, Flow Cytometry, Functional Assay, ELISA

PRODUCT SYNONYMS

MDH2; Polyclonal Antibody; Anti-MDH2 Antibody Picoband; anti-MDH2 antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Isotype	Rabbit IgG
Purity/Purification	Immunogen affinity purified.
Form/Format	Lyophilized Each vial contains 4 mg Trehalose, 0.9 mg NaCl, 0.2 mg Na ₂ HPO ₄ .
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml. (varies by lot)
Applicable Applications for anti-MDH2 antibody	WB (Western Blot), IHC (Immunohistochemistry), ICC (Immunocytochemistry), IF (Immunofluorescence), FCM/FACS (Flow Cytometry), ELISA
Application Notes	WB: 0.1-0.25 ug/ml, Human, Mouse, Rat IHC-P: 1-2 ug/ml, Human, Mouse, Rat ICC/IF: 5 ug/ml, Human FC/FACS: 1-3 ug/1x10 ⁶ cells, Human Direct ELISA: 0.1-0.5 ug/ml, Human Tested Species: In-house tested species with positive results. Enhanced Chemiluminescent Kit with anti-Rabbit IgG for Western blot, and HRP Conjugated anti-Rabbit IgG Super Vision Assay Kit for IHC(P) and ICC.
Reconstitution	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Immunogen	E Coli-derived human MDH2 recombinant protein (Position: A9-L223).
Preparation and Storage	Store at -20 degree C for one year from date of receipt. After reconstitution, at 4 degree C for one month. It can also be aliquotted and stored frozen at -20 degree C for six months. Avoid repeated freezing and thawing.




FCM (Flow Cytometry)

Figure 12. Flow Cytometry analysis of MCF-7 cells using anti-MDH2 antibody (AAA19527).Overlay histogram showing MCF-7 cells stained with AAA19527 (Blue line). The cells were blocked with 10% normal goat serum. And then incubated with rabbit anti-MDH2 Antibody (AAAA19527, 1 ug/1x10⁶ cells) for 30 min at 20 degree C. DyLight488 conjugated goat anti-rabbit IgG (BA1127, 5-10 ug/1x10⁶ cells) was used as secondary antibody for 30 minutes at 20 degree C. Isotype control antibody (Green line) was rabbit IgG (1 ug/1x10⁶) used under the same conditions. Unlabelled sample (Red line) was also used as a control.




IHC (Immunohistochemistry)

Figure 10. IHC analysis of MDH2 using anti-MDH2 antibody (AAA19527).MDH2 was detected in a paraffin-embedded section of rat brain tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 ug/ml rabbit anti-MDH2 Antibody (AAA19527) overnight at 4 degree C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37 degree C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC) with DAB as the chromogen.




IHC (Immunohistochemistry)

Figure 8. IHC analysis of MDH2 using anti-MDH2 antibody (AAA19527).MDH2 was detected in a paraffin-embedded section of mouse brain tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 ug/ml rabbit anti-MDH2 Antibody (AAA19527) overnight at 4 degree C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37 degree C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC) with DAB as the chromogen.




IF (Immunofluorescence)

Figure 11. IF analysis of MDH2 using anti-MDH2 antibody (AAA19527).MDH2 was detected in an immunocytochemical section of MCF-7 cells. Enzyme antigen retrieval was performed using IHC enzyme antigen retrieval reagent (AR0022) for 15 mins. The cells were blocked with 10% goat serum. And then incubated with 5 ug/mL rabbit anti-MDH2 Antibody (AAA19527) overnight at 4 degree C. DyLight594 Conjugated Goat Anti-Rabbit IgG (BA1142) was used as secondary antibody at 1:100 dilution and incubated for 30 minutes at 37 degree C. The section was counterstained with DAPI. Visualize using a fluorescence microscope and filter sets appropriate for the label used.



IHC (Immunohistochemistry)

Figure 9. IHC analysis of MDH2 using anti-MDH2 antibody (AAA19527).MDH2 was detected in a paraffin-embedded section of rat brain tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 ug/ml rabbit anti-MDH2 Antibody (AAA19527) overnight at 4 degree C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37 degree C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC) with DAB as the chromogen.



IHC (Immunohistochemistry)

Figure 7. IHC analysis of MDH2 using anti-MDH2 antibody (AAA19527).MDH2 was detected in a paraffin-embedded section of human thyroid papillary carcinoma tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 ug/ml rabbit anti-MDH2 Antibody (AAA19527) overnight at 4 degree C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37 degree C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC) with DAB as the chromogen.

ADDITIONAL INFORMATION

Related Product Information for anti-MDH2 antibody	Malate dehydrogenase, mitochondrial also known as malate dehydrogenase 2 is an enzyme that in humans is encoded by the MDH2 gene. Malate dehydrogenase catalyzes the reversible oxidation of malate to oxaloacetate, utilizing the NAD/NADH cofactor system in the citric acid cycle. The protein encoded by this gene is localized to the mitochondria and may play pivotal roles in the malate-aspartate shuttle that operates in the metabolic coordination between cytosol and mitochondria. Several transcript variants encoding different isoforms have been found for this gene.
Product Categories/Family for anti-MDH2 antibody	Primary Antibodies, Rabbit Polyclonal Antibodies

NCBI AND UNIPROT INFORMATION

NCBI GI #	538919685
NCBI GeneID	4191
NCBI Accession #	NP_001269332.1
NCBI GenBank Nucleotide #	NM_001282403.1
UniProt Accession #	P40926; O43682; A8K414; B2RE78; B4DE44; E9PDB2
NCBI Official Full Name	malate dehydrogenase, mitochondrial isoform 2
NCBI Official Synonym Full Names	malate dehydrogenase 2, NAD (mitochondrial)
NCBI Official Symbol	MDH2
NCBI Official Synonym Symbols	MDH; MOR1; M-MDH; MGC:3559
NCBI Protein Information	malate dehydrogenase, mitochondrial
UniProt Protein Name	Malate dehydrogenase, mitochondrial
UniProt Gene Name	MDH2
UniProt Entry Name	MDHM_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/mdh2-antibody-western-blot-immunohistochemistry-immunocytochemistry-immunofluorescence-flow-cytometry-elisa/19527>