

Rabbit anti-Human CD93 Polyclonal Antibody | anti-CD93 antibody

Anti-CD93 Antibody Picoband


CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA19531	Rabbit	Human	Immunohistochemistry, Flow Cytometry, Functional Assay

PRODUCT SYNONYMS

CD93; Polyclonal Antibody; Anti-CD93 Antibody Picoband; anti-CD93 antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	Rabbit IgG
Purity/Purification	Immunogen affinity purified.
Form/Format	Lyophilized Each vial contains 4 mg Trehalose, 0.9 mg NaCl, 0.2 mg Na ₂ HPO ₄ .
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml. (varies by lot)
Applicable Applications for anti-CD93 antibody	IHC (Immunohistochemistry), FCM/FACS (Flow Cytometry)
Application Notes	IHC-P: 1-2 ug/ml, Human FC/FACS: 1-3 ug/1x10 ⁶ cells, Human Tested Species: In-house tested species with positive results. HRP Conjugated anti-Rabbit IgG Super Vision Assay Kit for IHC(P).
Reconstitution	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Immunogen	A synthetic peptide corresponding to a sequence of human CD93 (QNAADSYSWVPERAESRAMENQ).
Preparation and Storage	Store at -20 degree C for one year from date of receipt. After reconstitution, at 4 degree C for one month. It can also be aliquotted and stored frozen at -20 degree C for six months. Avoid repeated freezing and thawing.




FCM (Flow Cytometry)

Figure 11. Flow Cytometry analysis of U937 cells using anti-CD93 antibody (AAA19531).Overlay histogram showing U937 cells stained with AAA19531 (Blue line). The cells were blocked with 10% normal goat serum. And then incubated with rabbit anti-CD93 Antibody (AAA19531, 1 ug/1x10⁶ cells) for 30 min at 20 degree C. DyLight488 conjugated goat anti-rabbit IgG was used as secondary antibody for 30 minutes at 20 degree C. Isotype control antibody (Green line) was rabbit IgG (1 ug/1x10⁶) used under the same conditions. Unlabelled sample (Red line) was also used as a control.




IHC (Immunohistochemistry)

Figure 9. IHC analysis of CD93 using anti-CD93 antibody (AAA19531).CD93 was detected in a paraffin-embedded section of human liver cancer tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 ug/ml rabbit anti-CD93 Antibody (AAA19531) overnight at 4 degree C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37 degree C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC) with DAB as the chromogen.




IHC (Immunohistochemistry)

Figure 7. IHC analysis of CD93 using anti-CD93 antibody (AAA19531).CD93 was detected in a paraffin-embedded section of human breast cancer tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 ug/ml rabbit anti-CD93 Antibody (AAA19531) overnight at 4 degree C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37 degree C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC) with DAB as the chromogen.




FCM (Flow Cytometry)

Figure 10. Flow Cytometry analysis of THP-1 cells using anti-CD93 antibody (AAA19531).Overlay histogram showing THP-1 cells stained with AAA19531 (Blue line). The cells were blocked with 10% normal goat serum. And then incubated with rabbit anti-CD93 Antibody (AAA19531, 1 ug/1x10⁶ cells) for 30 min at 20 degree C. DyLight488 conjugated goat anti-rabbit IgG was used as secondary antibody for 30 minutes at 20 degree C. Isotype control antibody (Green line) was rabbit IgG (1 ug/1x10⁶) used under the same conditions. Unlabelled sample (Red line) was also used as a control.



IHC (Immunohistochemistry)

Figure 8. IHC analysis of CD93 using anti-CD93 antibody (AAA19531).CD93 was detected in a paraffin-embedded section of human gall bladder adenocarcinoma tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 ug/ml rabbit anti-CD93 Antibody (AAA19531) overnight at 4 degree C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37 degree C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC) with DAB as the chromogen.



IHC (Immunohistochemistry)

Figure 6. IHC analysis of CD93 using anti-CD93 antibody (AAA19531).CD93 was detected in a paraffin-embedded section of human appendiceal adenocarcinoma tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 ug/ml rabbit anti-CD93 Antibody (AAA19531) overnight at 4 degree C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37 degree C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC) with DAB as the chromogen.

ADDITIONAL INFORMATION

Related Product Information for anti-CD93 antibody	CD93 (Cluster of Differentiation 93) is a protein that in humans is encoded by the CD93 gene. The protein encoded by this gene is a cell-surface glycoprotein and type I membrane protein that was originally identified as a myeloid cell-specific marker. The encoded protein was once thought to be a receptor for C1q, but now is thought to instead be involved in intercellular adhesion and in the clearance of apoptotic cells. The intracellular cytoplasmic tail of this protein has been found to interact with moesin, a protein known to play a role in linking transmembrane proteins to the cytoskeleton and in the remodelling of the cytoskeleton.
Product Categories/Family for anti-CD93 antibody	Primary Antibodies, Rabbit Polyclonal Antibodies

NCBI AND UNIPROT INFORMATION

NCBI GI #	88758613
NCBI GeneID	22918
NCBI Accession #	NP_036204.2
NCBI GenBank Nucleotide #	NM_012072.3
UniProt Accession #	Q9NPY3; O00274
Molecular Weight	68,560 Da
NCBI Official Full Name	complement component C1q receptor
NCBI Official Synonym Full Names	CD93 molecule
NCBI Official Symbol	CD93
NCBI Official Synonym Symbols	C1QR1; C1qRP; CDw93; ECSM3; MXRA4; C1qR(P); dJ737E23.1
NCBI Protein Information	complement component C1q receptor; C1qR; CD93 antigen; C1q receptor 1; C1q/MBL/SPA receptor; matrix-remodelling associated 4; matrix-modeling-associated protein 4; complement component 1 q subcomponent receptor 1; complement component 1, q subcomponent,
UniProt Protein Name	Complement component C1q receptor
UniProt Gene Name	CD93
UniProt Synonym Gene Names	C1QR1; MXRA4; C1qR; C1qR(p); C1qRp
UniProt Entry Name	C1QR1_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/cd93-antibody-immunohistochemistry-flow-cytometry-elisa/19531>