

Mouse HGS Monoclonal Antibody | anti-HGS antibody

Anti-HGS Antibody Picoband (monoclonal, 4B7E2)

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA19679	Mouse	Human, Mouse, Rat	Flow Cytometry, Functional Assay, Immunofluorescence, Immunohistochemistry, Immunocytochemistry, Western Blot

PRODUCT SYNONYMS

HGS; Monoclonal Antibody; Anti-HGS Antibody Picoband (monoclonal; 4B7E2); anti-HGS antibody


PRODUCT SPECIFICATIONS

Host	Mouse
Reactivity	Human, Mouse, Rat
Clonality	Monoclonal
Isotype	Mouse IgG2a
Clone Number	4B7E2
Purity/Purification	Immunogen affinity purified.
Form/Format	Lyophilized Each vial contains 4 mg Trehalose, 0.9 mg NaCl and 0.2 mg Na ₂ HPO ₄ .
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml. (varies by lot)
Applicable Applications for anti-HGS antibody	FCM/FACS (Flow Cytometry), IF (Immunofluorescence), IHC (Immunohistochemistry), ICC (Immunocytochemistry), WB (Western Blot)
Application Notes	WB: 0.25-0.5 ug/ml, Human, Mouse, Rat IHC-P: 2-5 ug/ml, Human, Mouse ICC/IF: 5 ug/ml, Human FC/FACS: 1-3 ug/1x10 ⁶ cells, Human Tested Species: In-house tested species with positive results. Enhanced Chemiluminescent Kit with anti-Mouse IgG for Western blot, and HRP Conjugated anti-Mouse IgG Super Vision Assay Kit for IHC(P) and ICC.
Reconstitution	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Immunogen	E Coli-derived human HGS recombinant protein (Position: R3-D777).
Preparation and Storage	Store at -20 degree C for one year from date of receipt. After reconstitution, at 4 degree C for one month. It can also be aliquotted and stored frozen at -20 degree C for six months. Avoid repeated freezing and thawing.




FCM (Flow Cytometry)

Figure 10. Flow Cytometry analysis of Caco-2 cells using anti-HGS antibody (AAA19679). Overlay histogram showing Caco-2 cells stained with AAA19679 (Blue line). The cells were blocked with 10% normal goat serum. And then incubated with mouse anti-HGS Antibody (AAA19679, 1 ug/1x10⁶ cells) for 30 min at 20 degree C. DyLight488 conjugated goat anti-mouse IgG was used as secondary antibody for 30 minutes at 20 degree C. Isotype control antibody (Green line) was mouse IgG (1 ug/1x10⁶) used under the same conditions. Unlabelled sample (Red line) was also used as a control.




IF (Immunofluorescence)

Figure 9. IF analysis of HGS using anti-HGS antibody (AAA19679). HGS was detected in an immunocytochemical section of U87 cells. Enzyme antigen retrieval was performed using IHC enzyme antigen retrieval reagent (AR0022) for 15 mins. The cells were blocked with 10% goat serum. And then incubated with 5 ug/mL mouse anti-HGS Antibody (AAA19679) overnight at 4 degree C. DyLight488 Conjugated Goat Anti-Mouse IgG was used as secondary antibody at 1:100 dilution and incubated for 30 minutes at 37 degree C. The section was counterstained with DAPI. Visualize using a fluorescence microscope and filter sets appropriate for the label used.



IHC (Immunohistochemistry)

Figure 8. IHC analysis of HGS using anti-HGS antibody (AAA19679). HGS was detected in a paraffin-embedded section of mouse testis tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 ug/ml mouse anti-HGS Antibody (AAA19679) overnight at 4 degree C. Peroxidase Conjugated Goat Anti-mouse IgG was used as secondary antibody and incubated for 30 minutes at 37 degree C. The tissue section was developed using HRP Conjugated Mouse IgG Super Vision Assay Kit with DAB as the chromogen.




IHC (Immunohistochemistry)

Figure 7. IHC analysis of HGS using anti-HGS antibody (AAA19679). HGS was detected in a paraffin-embedded section of human ovarian cancer tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 ug/ml mouse anti-HGS Antibody (AAA19679) overnight at 4 degree C. Peroxidase Conjugated Goat Anti-mouse IgG was used as secondary antibody and incubated for 30 minutes at 37 degree C. The tissue section was developed using HRP Conjugated Mouse IgG Super Vision Assay Kit with DAB as the chromogen.



IHC (Immunohistochemistry)

Figure 6. IHC analysis of HGS using anti-HGS antibody (AAA19679). HGS was detected in a paraffin-embedded section of human laryngeal squamous cell carcinomas tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 ug/ml mouse anti-HGS Antibody (AAA19679) overnight at 4 degree C. Peroxidase Conjugated Goat Anti-mouse IgG was used as secondary antibody and incubated for 30 minutes at 37 degree C. The tissue section was developed using HRP Conjugated Mouse IgG Super Vision Assay Kit with DAB as the chromogen.



IHC (Immunohistochemistry)

Figure 5. IHC analysis of HGS using anti-HGS antibody (AAA19679). HGS was detected in a paraffin-embedded section of human endometrial cancer tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 ug/ml mouse anti-HGS Antibody (AAA19679) overnight at 4 degree C. Peroxidase Conjugated Goat Anti-mouse IgG was used as secondary antibody and incubated for 30 minutes at 37 degree C. The tissue section was developed using HRP Conjugated Mouse IgG Super Vision Assay Kit with DAB as the chromogen.

ADDITIONAL INFORMATION

Related Product Information for anti-HGS antibody	Hepatocyte growth factor-regulated tyrosine kinase substrate is an enzyme that in humans is encoded by the HGS gene. It is mapped to 17q25.3. The protein encoded by this gene regulates endosomal sorting and plays a critical role in the recycling and degradation of membrane receptors. The encoded protein sorts monoubiquitinated membrane proteins into the multivesicular body, targeting these proteins for lysosome-dependent degradation.
Product Categories/Family for anti-HGS antibody	Primary Antibodies, Mouse Monoclonal Antibodies

NCBI AND UNIPROT INFORMATION

NCBI GI #	4758528
NCBI GeneID	9146
NCBI Accession #	NP_004703.1
NCBI GenBank Nucleotide #	NM_004712.4
UniProt Accession #	O14964; Q9NR36

Molecular Weight	86,192 Da
NCBI Official Full Name	hepatocyte growth factor-regulated tyrosine kinase substrate
NCBI Official Synonym Full Names	hepatocyte growth factor-regulated tyrosine kinase substrate
NCBI Official Symbol	HGS
NCBI Official Synonym Symbols	HRS
NCBI Protein Information	hepatocyte growth factor-regulated tyrosine kinase substrate; protein pp110; human growth factor-regulated tyrosine kinase substrate
UniProt Protein Name	Hepatocyte growth factor-regulated tyrosine kinase substrate
UniProt Gene Name	HGS
UniProt Synonym Gene Names	HRS
UniProt Entry Name	HGS_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/monoclonal/hgs-antibody-clone-4b7e2-flow-cytometry-immunofluorescence-immunohistochemistry-immunocytochemistry-western-blot-elisa/19679>