

Mouse MCM6 Monoclonal Antibody | anti-MCM6 antibody

Anti-MCM6 Antibody Picoband (monoclonal, 3F13C4)

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA19704	Mouse	Human, Mouse, Rat	Western Blot, Immunohistochemistry, Immunocytochemistry, Immunofluorescence, Flow Cytometry, Functional Assay

PRODUCT SYNONYMS

MCM6; Monoclonal Antibody; Anti-MCM6 Antibody Picoband (monoclonal; 3F13C4); anti-MCM6 antibody

PRODUCT SPECIFICATIONS

Host	Mouse
Reactivity	Human, Mouse, Rat
Clonality	Monoclonal
Isotype	Mouse IgG1
Clone Number	3F13C4
Purity/Purification	Immunogen affinity purified.
Form/Format	Lyophilized Each vial contains 4 mg Trehalose, 0.9 mg NaCl and 0.2 mg Na ₂ HPO ₄ .
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml. (varies by lot)
Applicable Applications for anti-MCM6 antibody	WB (Western Blot), IHC (Immunohistochemistry), ICC (Immunocytochemistry), IF (Immunofluorescence), FCM/FACS (Flow Cytometry)
Application Notes	WB: 0.25-0.5 ug/ml, Human, Mouse, Rat IHC-P: 2-5 ug/ml, Human, Mouse, Rat ICC/IF: 5 ug/ml, Human FC/FACS: 1-3 ug/1x10 ⁶ cells, Human Tested Species: In-house tested species with positive results. Enhanced Chemiluminescent Kit with anti-Mouse IgG for Western blot, and HRP Conjugated anti-Mouse IgG Super Vision Assay Kit for IHC(P) and ICC.
Reconstitution	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Immunogen	E Coli-derived human MCM6 recombinant protein (Position: Q14-D821).
Preparation and Storage	Store at -20 degree C for one year from date of receipt. After reconstitution, at 4 degree C for one month. It can also be aliquotted and stored frozen at -20 degree C for six months. Avoid repeated freezing and thawing.



FCM (Flow Cytometry)

Figure 11. Flow Cytometry analysis of HL-60 cells using anti-MCM6 antibody (AAA19704). Overlay histogram showing HL-60 cells stained with AAA19704 (Blue line). The cells were blocked with 10% normal goat serum. And then incubated with mouse anti-MCM6 Antibody (AAA19704, 1 ug/1x10⁶ cells) for 30 min at 20 degree C. DyLight488 conjugated goat anti-mouse IgG was used as secondary antibody for 30 minutes at 20 degree C. Isotype control antibody (Green line) was mouse IgG (1 ug/1x10⁶) used under the same conditions. Unlabelled sample (Red line) was also used as a control.



IF (Immunofluorescence)

Figure 10. IF analysis of MCM6 using anti-MCM6 antibody (AAA19704). MCM6 was detected in an immunocytochemical section of CACO-2 cells. Enzyme antigen retrieval was performed using IHC enzyme antigen retrieval reagent (AR0022) for 15 mins. The cells were blocked with 10% goat serum. And then incubated with 5 ug/mL mouse anti-MCM6 Antibody (AAA19704) overnight at 4 degree C. DyLight488 Conjugated Goat Anti-Mouse IgG was used as secondary antibody at 1:100 dilution and incubated for 30 minutes at 37 degree C. The section was counterstained with DAPI. Visualize using a fluorescence microscope and filter sets appropriate for the label used.



IHC (Immunohistochemistry)

Figure 9. IHC analysis of MCM6 using anti-MCM6 antibody (AAA19704). MCM6 was detected in a paraffin-embedded section of rat testis tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 ug/ml mouse anti-MCM6 Antibody (AAA19704) overnight at 4 degree C. Biotinylated goat anti-mouse IgG was used as secondary antibody and incubated for 30 minutes at 37 degree C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC) (#SA1021) with DAB as the chromogen.



IHC (Immunohistochemistry)

Figure 8. IHC analysis of MCM6 using anti-MCM6 antibody (AAA19704). MCM6 was detected in a paraffin-embedded section of mouse testis tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 ug/ml mouse anti-MCM6 Antibody (AAA19704) overnight at 4 degree C. Biotinylated goat anti-mouse IgG was used as secondary antibody and incubated for 30 minutes at 37 degree C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC) (#SA1021) with DAB as the chromogen.



IHC (Immunohistochemistry)

Figure 7. IHC analysis of MCM6 using anti-MCM6 antibody (AAA19704). MCM6 was detected in a paraffin-embedded section of human rectal moderately differentiate adenocarcinoma tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 ug/ml mouse anti-MCM6 Antibody (AAA19704) overnight at 4 degree C. Biotinylated goat anti-mouse IgG was used as secondary antibody and incubated for 30 minutes at 37 degree C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC) (#SA1021) with DAB as the chromogen.



IHC (Immunohistochemistry)

Figure 6. IHC analysis of MCM6 using anti-MCM6 antibody (AAA19704). MCM6 was detected in a paraffin-embedded section of human metaplasia of squamous cells of the renal pelvis tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 ug/ml mouse anti-MCM6 Antibody (AAA19704) overnight at 4 degree C. Biotinylated goat anti-mouse IgG was used as secondary antibody and incubated for 30 minutes at 37 degree C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC) (#SA1021) with DAB as the chromogen.

ADDITIONAL INFORMATION

Related Product Information for anti-MCM6 antibody	MCM6(Minichromosome maintenance, s. pombe, homolog of, 6) is a protein that in humans is encoded by the MCM6 gene. MCM6 is one of the highly conserved mini-chromosome maintenance proteins (MCM) that are essential for the initiation of eukaryotic genome replication. The MCM genes were originally identified in yeast defective in minichromosome maintenance and have since been shown to play roles in the progression of the cell cycle; many are cell division control genes. The MCM6 gene is mapped on 2q21.3. Mcm 6 has recently been shown to interact strongly Cdt1 at defined residues, by mutating these target residues Wei et al. observed lack of Cdt1 recruitment of Mcm2-7 to the pre-RC. An approximately 200-kb region surrounding the C/T(-13910) polymorphism in MCM6 intron 13 functioned as an enhancer of the lactase gene promoter in intestinal cell culture.
Product Categories/Family for anti-MCM6 antibody	Primary Antibodies, Mouse Monoclonal Antibodies

NCBI AND UNIPROT INFORMATION

NCBI GI #	2497824
NCBI GeneID	4175
UniProt Accession #	Q14566; Q13504; Q99859; B2R6H2

Molecular Weight	821
NCBI Official Full Name	DNA replication licensing factor MCM6
NCBI Official Synonym Full Names	minichromosome maintenance complex component 6
NCBI Official Symbol	MCM6
NCBI Official Synonym Symbols	Mis5; P105MCM; MCG40308
NCBI Protein Information	DNA replication licensing factor MCM6; minichromosome maintenance deficient (mis5, S. pombe) 6; MCM6 minichromosome maintenance deficient 6 (MIS5 homolog, S. pombe)
UniProt Protein Name	DNA replication licensing factor MCM6
UniProt Gene Name	MCM6
UniProt Entry Name	MCM6_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/monoclonal/mcm6-antibody-clone-3f13c4-western-blot-immunohistochemistry-immunocytochemistry-immunofluorescence-flow-cytometry-elisa/19704>