

Mouse anti-Human GSDMD Monoclonal Antibody | anti-GSDMD antibody

Anti-GSDMD Antibody Picoband (monoclonal, 6D11)


CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA19705	Mouse	Human	Western Blot, Immunohistochemistry, Immunocytochemistry, Immunofluorescence, Flow Cytometry, Functional Assay

PRODUCT SYNONYMS

GSDMD; Monoclonal Antibody; Anti-GSDMD Antibody Picoband (monoclonal; 6D11); anti-GSDMD antibody


PRODUCT SPECIFICATIONS

Host	Mouse
Reactivity	Human
Clonality	Monoclonal
Isotype	Mouse IgG2b
Clone Number	6D11
Purity/Purification	Immunogen affinity purified.
Form/Format	Lyophilized Each vial contains 4 mg Trehalose, 0.9 mg NaCl and 0.2 mg Na ₂ HPO ₄ .
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml. (varies by lot)
Applicable Applications for anti-GSDMD antibody	WB (Western Blot), IHC (Immunohistochemistry), ICC (Immunocytochemistry), IF (Immunofluorescence), FCM/FACS (Flow Cytometry)
Application Notes	WB: 0.25-0.5 ug/ml, Human IHC-P: 2-5 ug/ml, Human ICC/IF: 5 ug/ml, Human FC/FACS: 1-3 ug/1x10 ⁶ cells, Human Tested Species: In-house tested species with positive results. Enhanced Chemiluminescent Kit with anti-Mouse IgG for Western blot, and HRP Conjugated anti-Mouse IgG Super Vision Assay Kit for IHC(P) and ICC.
Reconstitution	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Immunogen	E Coli-derived human GSDMD recombinant protein (Position: M1-H484).
Preparation and Storage	Store at -20 degree C for one year from date of receipt. After reconstitution, at 4 degree C for one month. It can also be aliquotted and stored frozen at -20 degree C for six months. Avoid repeated freezing and thawing.




FCM (Flow Cytometry)

Figure 6. Flow Cytometry analysis of Jurkat cells using anti-GSDMD antibody (AAA19705).Overlay histogram showing Jurkat cells stained with AAA19705 (Blue line). The cells were blocked with 10% normal goat serum. And then incubated with mouse anti-GSDMD Antibody (AAA19705, 1 ug/1x10⁶ cells) for 30 min at 20 degree C. DyLight488 conjugated goat anti-mouse IgG was used as secondary antibody for 30 minutes at 20 degree C. Isotype control antibody (Green line) was mouse IgG (1 ug/1x10⁶) used under the same conditions. Unlabelled sample (Red line) was also used as a control.




IHC (Immunohistochemistry)

Figure 4. IHC analysis of GSDMD using anti-GSDMD antibody (AAA19705).GSDMD was detected in a paraffin-embedded section of human esophageal squamous carcinoma tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 ug/ml mouse anti-GSDMD Antibody (AAA19705) overnight at 4 degree C. Biotinylated goat anti-mouse IgG was used as secondary antibody and incubated for 30 minutes at 37 degree C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC) (#SA1021) with DAB as the chromogen.




IF (Immunofluorescence)

Figure 5. IF analysis of GSDMD using anti-GSDMD antibody (AAA19705).GSDMD was detected in an immunocytochemical section of PC-3 cells. Enzyme antigen retrieval was performed using IHC enzyme antigen retrieval reagent (AR0022) for 15 mins. The cells were blocked with 10% goat serum. And then incubated with 5 ug/mL mouse anti-GSDMD Antibody (AAA19705) overnight at 4 degree C. DyLight488 Conjugated Goat Anti-Mouse IgG was used as secondary antibody at 1:100 dilution and incubated for 30 minutes at 37 degree C. The section was counterstained with DAPI. Visualize using a fluorescence microscope and filter sets appropriate for the label used.




IHC (Immunohistochemistry)

Figure 3. IHC analysis of GSDMD using anti-GSDMD antibody (AAA19705).GSDMD was detected in a paraffin-embedded section of human cervical intraepithelial neoplasia tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 ug/ml mouse anti-GSDMD Antibody (AAA19705) overnight at 4 degree C. Biotinylated goat anti-mouse IgG was used as secondary antibody and incubated for 30 minutes at 37 degree C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC) (#SA1021) with DAB as the chromogen.



IHC (Immunohistochemistry)

Figure 2. IHC analysis of GSDMD using anti-GSDMD antibody (AAA19705). GSDMD was detected in a paraffin-embedded section of human cervical cancer tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 ug/ml mouse anti-GSDMD Antibody (AAA19705) overnight at 4 degree C. Biotinylated goat anti-mouse IgG was used as secondary antibody and incubated for 30 minutes at 37 degree C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC) (#SA1021) with DAB as the chromogen.



WB (Western Blot)

Figure 1. Western blot analysis of GSDMD using anti-GSDMD antibody (AAA19705). Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel)/90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions. Lane 1: human THP-1 whole cell lysates, Lane 2: human SW620 whole cell lysates, Lane 3: human Jurkat whole cell lysates, Lane 4: human U87 whole cell lysates, Lane 5: human PC-3 whole cell lysates. After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with mouse anti-GSDMD antigen affinity purified monoclonal antibody (#AAA19705) at 0.5 ug/mL overnight at 4 degree C, then washed with TBS-0.1% Tween 3 times with 5 minutes each and probed with a goat anti-mouse IgG-HRP secondary antibody at a dilution of 1:10000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit with Tanon 5200 system. A specific band was detected for GSDMD at approximately 53 kDa. The expected band size for GSDMD is at 53 kDa.

ADDITIONAL INFORMATION

Related Product Information for anti-GSDMD antibody	Gasdermin D is a member of the gasdermin family. Members of this family appear to play a role in regulation of epithelial proliferation. Gasdermin D has been suggested to act as a tumor suppressor. Alternatively spliced transcript variants have been described.
Product Categories/Family for anti-GSDMD antibody	Primary Antibodies, Mouse Monoclonal Antibodies

NCBI AND UNIPROT INFORMATION

NCBI GI #	83320070
NCBI GeneID	79792
NCBI Accession #	NP_079012.3
NCBI GenBank Nucleotide #	NM_024736.6
UniProt Accession #	P57764; Q96Q98; D3DWJ9
Molecular Weight	52,801 Da
NCBI Official Full Name	gasdermin-D
NCBI Official Synonym Full Names	gasdermin D
NCBI Official Symbol	GSDMD
NCBI Official Synonym Symbols	DF5L; DFNA5L; FKSG10; GSDMDC1
NCBI Protein Information	gasdermin-D; gasdermin domain containing 1; gasdermin domain-containing protein 1
UniProt Protein Name	Gasdermin-D
UniProt Gene Name	GSDMD
UniProt Synonym Gene Names	DFNA5L; GSDMDC1

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/monoclonal/gsdmd-antibody-clone-6d11-western-blot-immunohistochemistry-immunocytochemistry-immunofluorescence-flow-cytometry-elisa/19705>