

Rabbit GCLC Polyclonal Antibody | anti-GCLC antibody

Anti-GCLC Antibody Picoband


CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA19759	Rabbit	Human, Mouse, Rat	ELISA, Flow Cytometry, Functional Assay, Immunofluorescence, Immunohistochemistry, Western Blot

PRODUCT SYNONYMS

GCLC; Polyclonal Antibody; Anti-GCLC Antibody Picoband; anti-GCLC antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Purity/Purification	Immunogen affinity purified.
Form/Format	Lyophilized. Each vial contains 4 mg Trehalose, 0.9 mg NaCl, 0.2 mg Na ₂ HPO ₄ .
Applicable Applications for anti-GCLC antibody	ELISA, FCM/FACS (Flow Cytometry), IF (Immunofluorescence), IHC (Immunohistochemistry), WB (Western Blot)
Application Notes	WB: 0.25-0.5ug/ml, Human, Mouse, Rat IHC: 2-5ug/ml, Human IF: 5ug/ml, Human FC/FACS (Fixed): 1-3ug/1x10 ⁶ cells, Human ELISA: 0.1-0.5ug/ml
Immunogen	E coli-derived human GCLC recombinant protein (Position: R24-H388). Human GCLC shares 94.2% and 93.7% amino acid (aa) sequence identity with mouse and rat GCLC, respectively.
Reconstitution	Adding 0.2 ml of distilled water will yield a concentration of 500ug/ml.
Preparation and Storage	At -20 degree C for one year from date of receipt. After reconstitution, at 4 degree C for one month. It can also be aliquotted and stored frozen at -20 degree C for six months.Avoid repeated freezing and thawing.



IF (Immunofluorescence)

Figure 8. IF analysis of GCLC using anti-GCLC antibody (AAA19759).GCLC was detected in a paraffin-embedded section of human liver cancer tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 5ug/mL rabbit anti-GCLC Antibody (AAA19759) overnight at 4 degree C. DyLight594 Conjugated Goat Anti-Rabbit IgG (BA1142) was used as secondary antibody at 1:500 dilution and incubated for 30 minutes at 37 degree C. The section was counterstained with DAPI. Visualize using a fluorescence microscope and filter sets appropriate for the label used.



FCM (Flow Cytometry)

Figure 7. Flow Cytometry analysis of MCF-7 cells using anti-GCLC antibody (AAA19759).Overlay histogram showing MCF-7 cells stained with AAA19759 (Blue line). To facilitate intracellular staining, cells were fixed with 4% paraformaldehyde and permeabilized with permeabilization buffer. The cells were blocked with 10% normal goat serum. And then incubated with rabbit anti-GCLC Antibody (AAA19759, 1ug/1x10⁶ cells) for 30 min at 20 degree C. DyLight488 conjugated goat anti-rabbit IgG was used as secondary antibody for 30 minutes at 20 degree C. Isotype control antibody (Green line) was rabbit IgG (1ug/1x10⁶) used under the same conditions. Unlabelled sample (Red line) was also used as a control.

NCBI Accession #	NP_001184044.1
NCBI GenBank Nucleotide #	NM_001197115.1
UniProt Accession #	P48506 ; Q14399
Molecular Weight	72,766 Da
NCBI Official Full Name	glutamate--cysteine ligase catalytic subunit isoform b
NCBI Official Synonym Full Names	glutamate-cysteine ligase, catalytic subunit
NCBI Official Symbol	GCLC
NCBI Official Synonym Symbols	GCL; GCS; GLCL; GLCLC
NCBI Protein Information	glutamate--cysteine ligase catalytic subunit; gamma-ECS; GCS heavy chain; gamma-glutamylcysteine synthetase
UniProt Protein Name	Glutamate--cysteine ligase catalytic subunit
UniProt Gene Name	GCLC
UniProt Synonym Gene Names	GLCL; GLCLC
UniProt Entry Name	GSH1_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/gclc-antibody-flow-cytometry-immunofluorescence-immunohistochemistry-western-blot-elisa/19759>