

Rabbit anti-Human CD127/IL7R Polyclonal Antibody | anti-CD127/IL7R antibody

Anti-CD127/IL7R Antibody Picoband


CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA19771	Rabbit	Human	Western Blot, Immunohistochemistry, Flow Cytometry, Functional Assay, ELISA

PRODUCT SYNONYMS

CD127/IL7R; Polyclonal Antibody; Anti-CD127/IL7R Antibody Picoband; Serine/threonine-protein kinase AtPK1/AtPK6; 70 kDa ribosomal protein S6 kinase 1 antibody; KS6B1_HUMAN antibody; p70 alpha antibody; P70 beta 1 antibody; p70 ribosomal S6 kinase alpha antibody; p70 ribosomal S6 kinase beta 1 antibody; p70 S6 kinase alpha antibody; P70 S6 Kinase antibody; p70 S6 kinase alpha 1 antibody; p70 S6 kinase alpha 2 antibody; p70 S6K antibody; p70 S6K-alpha antibody; p70 S6KA antibody; p70(S6K) alpha antibody; p70(S6K)-alpha antibody; p70-alpha antibody; p70-S6K 1 antibody; p70-S6K antibody; P70S6K antibody; P70S6K1 antibody; p70S6Kb antibody; PS6K antibody; Ribosomal protein S6 kinase 70kDa polypeptide 1 antibody; Ribosomal protein S6 kinase beta 1 antibody; Ribosomal protein S6 kinase beta-1 antibody; Ribosomal protein S6 kinase I antibody; RPS6KB1 antibody; S6K antibody; S6K-beta-1 antibody; S6K1 antibody; Serine/threonine kinase 14 alpha antibody; Serine/threonine-protein kinase 14A antibody; STK14A antibody; anti-CD127/IL7R antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Isotype	IgG
Specificity	Expressed in all tissues.
Purity/Purification	Immunogen affinity purified.
Form/Format	Lyophilized. Each vial contains 4 mg Trehalose, 0.9 mg NaCl, 0.2 mg Na2HPO4.
Applicable Applications for anti-CD127/IL7R antibody	WB (Western Blot), IHC (Immunohistochemistry), FCM/FACS (Flow Cytometry), ELISA
Application Notes	WB: 0.25-0.5ug/ml, Human IHC: 1-2ug/ml, Human FC/FACS (Fixed): 1-3ug/1x10 ⁶ cells, Human ELISA: 0.1-0.5ug/ml
Immunogen	E coli-derived human CD127/IL7R recombinant protein (Position: A33-Q433). Human IL7R shares 63.1% amino acid (aa) sequence identity with mouse IL7R.
Subcellular Localization	Cytoplasm, Nucleus
Reconstitution	Adding 0.2 ml of distilled water will yield a concentration of 500ug/ml.
Cross Reactivity	No cross reactivity with other proteins.
Protein Function	Downstream effector of TOR signaling pathway involved in osmotic stress response. Could be involved in the control of plant growth and development. Phosphorylates the ribosomal proteins P14, P16 and S6. Functions as a repressor of cell proliferation and required for maintenance of chromosome stability and ploidy levels through the RBR1-E2F pathway.
Preparation and Storage	At -20 degree C for one year from date of receipt. After reconstitution, at 4 degree C for one month. It can also be aliquotted and stored frozen at -20 degree C for six months.Avoid repeated freezing and thawing.



FCM (Flow Cytometry)

Figure 6. Flow Cytometry analysis of JK cells using anti-CD127/IL7R antibody (AAA19771). Overlay histogram showing JK cells stained with AAA19771 (Blue line). The cells were fixed with 4% paraformaldehyde and blocked with 10% normal goat serum. And then incubated with rabbit anti-CD127/IL7R Antibody (AAA19771, 1ug/1x10⁶ cells) for 30 min at 20 degree C. Dylight488 conjugated goat anti-rabbit IgG was used as secondary antibody for 30 minutes at 20 degree C. Isotype control antibody (Green line) was rabbit IgG (1ug/1x10⁶) used under the same conditions. Unlabelled sample (Red line) was also used as a control.



IHC (Immunohistochemistry)

Figure 5. IHC analysis of CD127/IL7R using anti-CD127/IL7R antibody (AAA19771). CD127/IL7R was detected in a paraffin-embedded section of human colon tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2ug/ml rabbit anti-CD127/IL7R Antibody (AAA19771) overnight at 4 degree C. Peroxidase Conjugated Goat Anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37 degree C. The tissue section was developed using HRP Conjugated Rabbit IgG Super Vision Assay Kit (epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2ug/ml rabbit anti-CD127/IL7R Antibody (AAA19771) overnight at 4 degree C. Peroxidase Conjugated Goat Anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37 degree C. The tissue section was developed using HRP Conjugated Rabbit IgG Super Vision Assay Kit (epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2ug/ml rabbit anti-CD127/IL7R Antibody (AAA19771) overnight at 4 degree C. Peroxidase Conjugated Goat Anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37 degree C. The tissue section was developed using HRP Conjugated Rabbit IgG Super Vision Assay Kit (epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2ug/ml rabbit anti-CD127/IL7R Antibody (AAA19771) overnight at 4 degree C. Peroxidase Conjugated Goat Anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37 degree C. The tissue section was developed using HRP Conjugated Rabbit IgG Super Vision Assay Kit (epitope retrieval solution).

The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2ug/ml rabbit anti-CD127/IL7R Antibody (AAA19771) overnight at 4 degree C. Peroxidase Conjugated Goat Anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37 degree C. The tissue section was developed using HRP Conjugated Rabbit IgG Super Vision Assay Kit (epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2ug/ml rabbit anti-CD127/IL7R Antibody (AAA19771) overnight at 4 degree C. Peroxidase Conjugated Goat Anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37 degree C. The tissue section was developed using HRP Conjugated Rabbit IgG Super Vision Assay Kit (epitope retrieval solution).

Lane 2: human K562 whole cell lysates, Lane 3: human Jurkat whole cell lysates. After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-CD127/IL7R antigen affinity purified polyclonal antibody (#AAA19771) at 0.5ug/mL overnight at 4 degree C, then washed with TBS+0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit with Tanon 5200 system. A specific band was detected for CD127/IL7R at approximately 170 kDa. The expected band size for CD127/IL7R is at 52 kDa.

ADDITIONAL INFORMATION

Related Product Information for anti-CD127/IL7R antibody	The brand Picoband indicates this is a premium antibody that guarantees superior quality, high affinity, and strong signals with minimal background in Western blot applications. Only our best-performing antibodies are designated as Picoband, ensuring unmatched performance.
Product Categories/Family for anti-CD127/IL7R antibody	Primary Antibodies ; Signal Transduction ; Signaling Pathway ; Small G Proteins
References	1. "Entrez Gene: IL7R interleukin 7 receptor".2. Kroemer RT, Richards WG (December 1996). "Homology modeling study of the human interleukin-7 receptor complex".Protein Eng. 9 (12): 1135-42.3. O'Doherty C, Alloza I, Rooney M, Vandenbroeck K (November 2009). "IL7RA polymorphisms and chronic inflammatory arthropathies". Tissue Antigens 74 (5): 429-31.

NCBI AND UNIPROT INFORMATION

NCBI GI #	28610151
-----------	----------

NCBI GeneID	3575
NCBI Accession #	NP_002176.2
NCBI GenBank Nucleotide #	NM_002185.3
UniProt Accession #	P16871; Q05CU8; Q6NSP4; Q6NWM0; Q6NWM1; Q6NWM2; Q6NWM3; Q6SV45; Q9UPC1; B2RCS6; B4DVT1
Molecular Weight	51,581 Da
NCBI Official Full Name	interleukin-7 receptor subunit alpha
NCBI Official Synonym Full Names	interleukin 7 receptor
NCBI Official Symbol	IL7R
NCBI Official Synonym Symbols	ILRA; CD127; IL7RA; CDW127; IL-7R-alpha
NCBI Protein Information	interleukin-7 receptor subunit alpha; IL-7RA; CD127 antigen; IL-7R subunit alpha; IL-7 receptor subunit alpha; interleukin 7 receptor alpha chain; interleukin 7 receptor isoform H5-6
UniProt Protein Name	Interleukin-7 receptor subunit alpha
UniProt Gene Name	IL7R
UniProt Synonym Gene Names	IL-7 receptor subunit alpha; IL-7R subunit alpha; IL-7R-alpha; IL-7RA
UniProt Entry Name	IL7RA_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/cd127-il7r-antibody-western-blot-immunohistochemistry-flow-cytometry-elisa/19771>