

Rabbit NXF1 Polyclonal Antibody | anti-NXF1 antibody

NXF1 antibody - N-terminal region

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA198726	Rabbit	Cow, Dog, Guinea Pig, Horse, Human, Mouse, Rabbit, Rat	Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

NXF1; Polyclonal Antibody; NXF1 antibody - N-terminal region; anti-NXF1 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Cow, Dog, Guinea Pig, Horse, Human, Mouse, Rabbit, Rat
Clonality	Polyclonal
Purity/Purification	Affinity Purified
Form/Format	Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.
Sequence	Synthetic peptide with the following region: RP NRRGDTWHDR DRIHVTVRRD RAPPERGGAG TSQDGTSKNW FKITIPYG
Sequence Length	619
Applicable Applications for anti-NXF1 antibody	IHC (Immunohistochemistry), WB (Western Blot)
Homology	Cow: 100%; Dog: 100%; Guinea Pig: 100%; Horse: 93%; Human: 100%; Mouse: 93%; Rabbit: 100%; Rat: 100%
Immunogen	The immunogen is a synthetic peptide directed towards the N terminal region of human NXF1
Preparation and Storage	For short term use, store at 2-8 degree C up to 1 week. For long term storage, store at -20 degree C in small aliquots to prevent freeze-thaw cycles.

IMAGES



WB (Western Blot)

WB Suggested Anti-NXF1 Antibody Titration: 0.2-1 ug/ml
Positive Control: HepG2 cell lysate
NXF1 is supported by BioGPS gene expression data to be expressed in HepG2



IHC (Immunohistochemistry)

Rabbit Anti-NXF1 Antibody
Formalin Fixed Paraffin Embedded Tissue: Human heart
Tissue
Observed Staining: Cytoplasmic, nucleus
Primary Antibody Concentration: N/A
Other Working Concentrations: 1:600
Secondary Antibody: Donkey anti-Rabbit-Cy3
Secondary Antibody Concentration: 1:200
Magnification: 20X
Exposure Time: 0.5 - 2.0 sec

ADDITIONAL INFORMATION

Related Product Information for anti-NXF1 antibody	This is a rabbit polyclonal antibody against NXF1. It was validated on Western Blot using a cell lysate as a positive control. Target Description: NXF1 is one member of a family of nuclear RNA export factor. Common domain features of this family are a noncanonical RNP-type RNA-binding domain (RBD), 4 leucine-rich repeats (LRRs), a nuclear transport factor 2 (NTF2)-like domain that allows heterodimerization with NTF2-related export protein-1 (NXT1), and a ubiquitin-associated domain that mediates interactions with nucleoporins. The LRRs and NTF2-like domains are required for export activity. NXF1 shuttles between the nucleus and the cytoplasm and binds in vivo to poly(A)+ RNA. NXF1 overcomes the mRNA export block caused by the presence of saturating amounts of CTE (constitutive transport element) RNA of type D retroviruses.This gene is one member of a family of nuclear RNA export factor genes. Common domain features of this family are a noncanonical RNP-type RNA-binding domain (RBD), 4 leucine-rich repeats (LRRs), a nuclear transport factor 2 (NTF2)-like domain that allows heterodimerization with NTF2-related export protein-1 (NXT1), and a ubiquitin-associated domain that mediates interactions with nucleoporins. The LRRs and NTF2-like domains are required for export activity. Alternative splicing seems to be a common mechanism in this gene family. The encoded protein of this gene shuttles between the nucleus and the cytoplasm and binds in vivo to poly(A)+ RNA. It is the vertebrate homologue of the yeast protein Mex67p. The encoded protein overcomes the mRNA export block caused by the presence of saturating amounts of CTE (constitutive transport element) RNA of type D retroviruses.This gene is one member of a family of nuclear RNA export factor genes. Common domain features of this family are a noncanonical RNP-type RNA-binding domain (RBD), 4 leucine-rich repeats (LRRs), a nuclear transport factor 2 (NTF2)-like domain that allows heterodimerization with NTF2-related export protein-1 (NXT1), and a ubiquitin-associated domain that mediates interactions with nucleoporins. The LRRs and NTF2-like domains are required for export activity. Alternative splicing seems to be a common mechanism in this gene family. The encoded protein of this gene shuttles between the nucleus and the cytoplasm and binds in vivo to poly(A)+ RNA. It is the vertebrate homologue of the yeast protein Mex67p. The encoded protein overcomes the mRNA export block caused by the presence of saturating amounts of CTE (constitutive transport element) RNA of type D retroviruses.
Product Categories/Family for anti-NXF1 antibody	Polyclonal; RNA Binding Proteins; Chromatin & Nuclear Signaling; E3 & Ubiquitin; DNA/RNA/Protein Interactions

NCBI AND UNIPROT INFORMATION

NCBI GI #	15487670
NCBI GeneID	10482
NCBI Accession #	NP_006353
NCBI GenBank Nucleotide #	NM_006362
UniProt Accession #	Q9UBU9
Molecular Weight	68kDa
NCBI Official Full Name	nuclear RNA export factor 1 isoform 1
NCBI Official Synonym Full Names	nuclear RNA export factor 1
NCBI Official Symbol	NXF1
NCBI Official Synonym Symbols	TAP; MEX67
NCBI Protein Information	nuclear RNA export factor 1
UniProt Protein Name	Nuclear RNA export factor 1
UniProt Gene Name	NXF1
UniProt Synonym Gene Names	TAP
UniProt Entry Name	NXF1_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

