

Rabbit MAT1A Polyclonal Antibody | anti-MAT1A antibody

MAT1A antibody - N-terminal region

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA198888	Rabbit	Cow, Dog, Guinea Pig, Horse, Human, Mouse, Rabbit, Rat, Zebrafish	Western Blot, Immunohistochemistry


PRODUCT SYNONYMS

MAT1A; Polyclonal Antibody; MAT1A antibody - N-terminal region; anti-MAT1A antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Cow, Dog, Guinea Pig, Horse, Human, Mouse, Rabbit, Rat, Zebrafish
Clonality	Polyclonal
Purity/Purification	Protein A purified
Form/Format	Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.
Sequence	Synthetic peptide with the following region: TS ESVGEGHPDK ICDQISDAVL DAHLKQDPNA KVACETVCKT GMVLLCGE
Sequence Length	395
Applicable Applications for anti-MAT1A antibody	WB (Western Blot), IHC (Immunohistochemistry)
Homology	Cow: 100%; Dog: 100%; Guinea Pig: 100%; Horse: 100%; Human: 100%; Mouse: 100%; Rabbit: 100%; Rat: 100%; Zebrafish: 93%
Immunogen	The immunogen is a synthetic peptide directed towards the N terminal region of human MAT1A
Preparation and Storage	For short term use, store at 2-8 degree C up to 1 week. For long term storage, store at -20 degree C in small aliquots to prevent freeze-thaw cycles.

IMAGES



WB (Western Blot)

WB Suggested Anti-MAT1A Antibody Titration: 1.25 µg/ml Positive Control: Jurkat cell lysate



IHC (Immunohistochemistry)

Rabbit Anti-MAT1A Antibody Paraffin Embedded Tissue: Human Kidney Cellular Data: Epithelial cells of renal tubule Antibody Concentration: 4.0-8.0 µg/ml Magnification: 400X

ADDITIONAL INFORMATION

Related Product Information for anti-MAT1A antibody	This is a rabbit polyclonal antibody against MAT1A. It was validated on Western Blot and immunohistochemistry Target Description: MAT1A catalyzes the formation of S-adenosylmethionine from methionine and ATP. Methionine adenosyltransferase deficiency is caused by recessive and dominant mutations, the latter identified in autosomal dominant persistant hypermethioninemia. This gene encodes methionine adenosyltransferase I (alpha isoform), which catalyzes the formation of S-adenosylmethionine from methionine and ATP. Methionine adenosyltransferase deficiency is caused by recessive and dominant mutations, the latter identified in autosomal dominant persistant hypermethioninemia.
Product Categories/Family for anti-MAT1A antibody	Polyclonal; Signal Proteins; Lymphocyte Signaling; Membrane Protein; Drugs and Drug Metabolism; Developmental Biology; Immunohistochemistry; Receptors; Methylation

NCBI AND UNIPROT INFORMATION

NCBI GI #	4557737
NCBI GeneID	4143
NCBI Accession #	NP_000420
NCBI GenBank Nucleotide #	NM_000429
UniProt Accession #	Q00266
Molecular Weight	43kDa
NCBI Official Full Name	S-adenosylmethionine synthase isoform type-1
NCBI Official Synonym Full Names	methionine adenosyltransferase 1A
NCBI Official Symbol	MAT1A
NCBI Official Synonym Symbols	MAT; SAMS; MATA1; SAMS1
NCBI Protein Information	S-adenosylmethionine synthase isoform type-1
UniProt Protein Name	S-adenosylmethionine synthase isoform type-1
UniProt Gene Name	MAT1A
UniProt Synonym Gene Names	AMS1; MATA1; AdoMet synthase 1; MAT 1; MAT-I/III
UniProt Entry Name	METK1_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/mat1a-antibody-western-blot-immunohistochemistry-elisa/198888>