

Rabbit UBE2D2 Polyclonal Antibody | anti-UBE2D2 antibody

UBE2D2 antibody - middle region

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA199208	Rabbit	Cow, Dog, Guinea Pig, Horse, Human, Mouse, Rabbit, Rat, Zebrafish	Western Blot, Immunohistochemistry


PRODUCT SYNONYMS

UBE2D2; Polyclonal Antibody; UBE2D2 antibody - middle region; anti-UBE2D2 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Cow, Dog, Guinea Pig, Horse, Human, Mouse, Rabbit, Rat, Zebrafish
Clonality	Polyclonal
Purity/Purification	Protein A purified
Form/Format	Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.
Sequence	Synthetic peptide with the following region: SQ WSPALTISKV LLSICSLLCD PNPDDPLVPE IARIYKTDRE KYNRIARE
Sequence Length	547
Applicable Applications for anti-UBE2D2 antibody	WB (Western Blot), IHC (Immunohistochemistry)
Homology	Cow: 100%; Dog: 100%; Guinea Pig: 100%; Horse: 100%; Human: 100%; Mouse: 100%; Rabbit: 100%; Rat: 100%; Zebrafish: 100%
Immunogen	The immunogen is a synthetic peptide directed towards the middle region of human UBE2D2
Preparation and Storage	For short term use, store at 2-8 degree C up to 1 week. For long term storage, store at -20 degree C in small aliquots to prevent freeze-thaw cycles.

IMAGES



WB (Western Blot)

WB Suggested Anti-UBE2D2 Antibody Titration: 1.25ug/ml Positive Control: Jurkat cell lysate



IHC (Immunohistochemistry)

Human Lung

ADDITIONAL INFORMATION

Related Product Information for anti-UBE2D2 antibody	This is a rabbit polyclonal antibody against UBE2D2. It was validated on Western Blot and immunohistochemistry Target Description: The modification of proteins with ubiquitin is an important cellular mechanism for targeting abnormal or short-lived proteins for degradation. Ubiquitination involves at least three classes of enzymes: ubiquitin-activating enzymes, or E1s, ubiquitin-conjugating enzymes, or E2s, and ubiquitin-protein ligases, or E3s. UBE2D2 is a member of the E2 ubiquitin-conjugating enzyme family. This enzyme functions in the ubiquitination of the tumor-suppressor protein p53, which is induced by an E3 ubiquitin-protein ligase.SLC38A4 is found predominantly in liver and transports both cationic and neutral amino acids. The transport of cationic amino acids by SLC38A4 is Na(+) and pH independent, while the transport of neutral amino acids is Na(+) and pH dependent (Hatanaka et al., 2001).[supplied by OMIM].
Product Categories/Family for anti-UBE2D2 antibody	Polyclonal; Membrane Protein; E3 & Ubiquitin; Immunohistochemistry

NCBI AND UNIPROT INFORMATION

NCBI GI #	4507775
NCBI GeneID	7322
NCBI Accession #	NP_003330
NCBI GenBank Nucleotide #	NM_018018
UniProt Accession #	P62837
Molecular Weight	60kDa
NCBI Official Full Name	ubiquitin-conjugating enzyme E2 D2 isoform 1
NCBI Official Synonym Full Names	ubiquitin conjugating enzyme E2 D2
NCBI Official Symbol	UBE2D2
NCBI Official Synonym Symbols	UBC4; PUBC1; UBCH4; UBC4/5; UBCH5B; E2(17)KB2
NCBI Protein Information	ubiquitin-conjugating enzyme E2 D2
UniProt Protein Name	Ubiquitin-conjugating enzyme E2 D2
UniProt Gene Name	UBE2D2
UniProt Synonym Gene Names	PUBC1; UBC4; UBC5B; UBCH4; UBCH5B
UniProt Entry Name	UB2D2_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/ube2d2-antibody-western-blot-immunohistochemistry-elisa/199208>