

Rabbit MELTF Polyclonal Antibody | anti-MELTF antibody

Anti-MELTF Antibody Picoband


CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA19980	Rabbit	Human, Mouse, Rat	Western Blot, Immunohistochemistry, ELISA

PRODUCT SYNONYMS

MELTF; Polyclonal Antibody; Anti-MELTF Antibody Picoband; anti-MELTF antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Purity/Purification	Immunogen affinity purified.
Form/Format	Lyophilized. Each vial contains 4 mg Trehalose, 0.9 mg NaCl, 0.2 mg Na ₂ HPO ₄ .
Applicable Applications for anti-MELTF antibody	WB (Western Blot), IHC (Immunohistochemistry), ELISA
Application Notes	WB: 0.25-0.5ug/ml, Human, Mouse, Rat IHC: 2-5ug/ml, Human, Mouse, Rat ELISA: 0.1-0.5ug/ml
Immunogen	E coli-derived human MFI2/MELTF recombinant protein (Position: A60-D647). Human MFI2/MELTF shares 86.7% amino acid (aa) sequence identity with mouse MFI2/MELTF.
Reconstitution	Adding 0.2 ml of distilled water will yield a concentration of 500ug/ml.
Preparation and Storage	At -20 degree C for one year from date of receipt. After reconstitution, at 4 degree C for one month. It can also be aliquotted and stored frozen at -20 degree C for six months.Avoid repeated freezing and thawing.



IHC (Immunohistochemistry)

Figure 8. IHC analysis of MELTF using anti-MELTF antibody (AAA19980).MELTF was detected in a paraffin-embedded section of rat testis tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2ug/ml rabbit anti-MELTF Antibody (AAA19980) overnight at 4 degree C. Peroxidase Conjugated Goat Anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37 degree C. The tissue section was developed using HRP Conjugated Rabbit IgG Super Vision Assay Kit (epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2ug/ml rabbit anti-MELTF Antibody (AAA19980) overnight at 4 degree C. Peroxidase Conjugated Goat Anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37 degree C. The tissue section was developed using HRP Conjugated Rabbit IgG Super Vision Assay Kit (epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2ug/ml rabbit anti-MELTF Antibody (AAA19980) overnight at 4 degree C. Peroxidase Conjugated Goat Anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37 degree C. The tissue section was developed using HRP Conjugated Rabbit IgG Super Vision Assay Kit (epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2ug/ml rabbit anti-MELTF Antibody (AAA19980) overnight at 4 degree C. Peroxidase Conjugated Goat Anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37 degree C. The tissue section was developed using HRP Conjugated Rabbit IgG Super Vision Assay Kit (epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2ug/ml rabbit anti-MELTF Antibody (AAA19980) overnight at 4 degree C. Peroxidase Conjugated Goat Anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37 degree C. The tissue section was developed using HRP Conjugated Rabbit IgG Super Vision Assay Kit (epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2ug/ml rabbit anti-MELTF Antibody (AAA19980) overnight at 4 degree C. Peroxidase Conjugated Goat Anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37 degree C. The tissue section was developed using HRP Conjugated Rabbit IgG Super Vision Assay Kit (Lane 2: rat C6 whole cell lysates,Lane 3: mouse skin tissue lysates.After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-MELTF antigen affinity purified polyclonal antibody (#AAA19980) at 0.5ug/mL overnight at 4 degree C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Lane 2: human A431 whole cell lysates,Lane 3: human A549 whole cell lysates,Lane 4: human PC-3 whole cell lysates.After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-MELTF antigen affinity purified polyclonal antibody (#AAA19980) at 0.5ug/mL overnight at 4 degree C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit with Tanon 5200 system. A specific band was detected for MELTF at approximately 90 kDa. The expected band size for MELTF is at 80 kDa.

ADDITIONAL INFORMATION

Related Product Information for anti-MELTF antibody	The brand Picoband indicates this is a premium antibody that guarantees superior quality, high affinity, and strong signals with minimal background in Western blot applications. Only our best-performing antibodies are designated as Picoband, ensuring unmatched performance.
Product Categories/Family for anti-MELTF antibody	Primary Antibodies
References	1. Le Beau, M. M., Diaz, M. O., Plowman, G. D., Brown, J. P., Rowley, J. D. Chromosomal sublocalization of the human p97 melanoma antigen. Hum. Genet. 72: 294-296, 1986.2. Plowman, G. D., Brown, J. P., Enns, C. A., Schroder, J., Nikinmaa, B., Sussman, H. H., Hellstrom, K. E., Hellstrom, I. Assignment of the gene for human melanoma-associated antigen p97 to chromosome 3. Nature 303: 70-72, 1983.3. Rose, T. M., Plowman, G. D., Teplow, D. B., Dreyer, W. J., Hellstrom, K. E., Brown, J. P. Primary structure of the human melanoma-associated antigen p97 (melanotransferrin) deduced from the mRNA sequence. Proc. Nat. Acad. Sci. 83: 1261-1265, 1986.

NCBI AND UNIPROT INFORMATION

NCBI GI #	134244281
NCBI GeneID	4241
NCBI Accession #	NP_005920.2
NCBI GenBank Nucleotide #	NM_005929.5
UniProt Accession #	P08582; Q9BQE2
Molecular Weight	80,215 Da
NCBI Official Full Name	melanotransferrin isoform 1
NCBI Official Synonym Full Names	antigen p97 (melanoma associated) identified by monoclonal antibodies 133.2 and 96.5
NCBI Official Symbol	MFI2
NCBI Official Synonym Symbols	MTF1; CD228; MAP97
NCBI Protein Information	melanotransferrin; membrane-bound transferrin-like protein; melanoma-associated antigen p97, isoform 2
UniProt Protein Name	Melanotransferrin
UniProt Gene Name	MFI2
UniProt Synonym Gene Names	MAP97
UniProt Entry Name	TRFM_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/meltf-antibody-western-blot-immunohistochemistry-elisa/19980>