

Rabbit anti-Human Interleukin 35 (IL35) Polyclonal Antibody | anti-IL35 antibody

Polyclonal Antibody to Interleukin 35 (IL35)

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA20105	Rabbit	Human	Western Blot, Immunohistochemistry, Immunocytochemistry, Immunoprecipitation


PRODUCT SYNONYMS

Interleukin 35 (IL35); Polyclonal Antibody; Polyclonal Antibody to Interleukin 35 (IL35); anti-IL35 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Purity/Purification	Antigen-specific affinity chromatography followed by Protein A affinity chromatography
Form/Format	Supplied as solution form in 0.01M PBS, pH7.4, containing 0.05% Proclin-300, 50% glycerol.
Concentration	0.5mg/mL (varies by lot)
Applicable Applications for anti-IL35 antibody	WB (Western Blot), IHC (Immunohistochemistry), ICC (Immunocytochemistry), IP (Immunoprecipitation)
Application Notes	Western blotting: 0.01-3ug/mL; Immunohistochemistry: 5-30ug/mL; Immunocytochemistry: 5-30ug/mL; Optimal working dilutions must be determined by end user.
Source	Polyclonal antibody preparation
Traits	Liquid
Immunogen	Recombinant IL35 (The IL35 complex composed of IL27B subunit (Arg21-Lys229) and the mature form of human IL12A (Arg23-Ser219) linked by a polypeptide linker (GGGGS)3) expressed in E.coli .
Preparation and Storage	Storage: Avoid repeated freeze/thaw cycles. Store at 4 degree C for frequent use. Aliquot and store at -20 degree C for 24 months. Stability Test: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37 degree C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

IMAGES



IHC (Immunohistochemistry)

DAB staining on IHC-P; Samples: Human Liver Tissue; Primary Ab: 30ug/ml Rabbit Anti-Human IL35 Antibody; Second Ab: 2ug/mL HRP-Linked Caprine Anti-Rabbit IgG Polyclonal Antibody (Catalog: AAA20105)



IHC (Immunohistochemistry)

DAB staining on IHC-P; Samples: Human Prostate cancer Tissue; Primary Ab: 30ug/ml Rabbit Anti-Human IL35 Antibody; Second Ab: 2ug/mL HRP-Linked Caprine Anti-Rabbit IgG Polyclonal Antibody (Catalog: AAA20105)

**IHC (Immunohistochemistry)**


DAB staining on IHC-P; Samples: Human Colorectal cancer Tissue; Primary Ab: 30ug/ml Rabbit Anti-Human IL35 Antibody; Second Ab: 2ug/mL HRP-Linked Caprine Anti-Rabbit IgG Polyclonal Antibody (Catalog:

**IHC (Immunohistochemistry)**

DAB staining on IHC-P; Samples: Human Stomach Tissue; Primary Ab: 30ug/ml Rabbit Anti-Human IL35 Antibody; Second Ab: 2ug/mL HRP-Linked Caprine Anti-Rabbit IgG Polyclonal Antibody (Catalog:

**IHC (Immunohistochemistry)**

DAB staining on IHC-P; Samples: Human Glioma Tissue; Primary Ab: 30ug/ml Rabbit Anti-Human IL35 Antibody; Second Ab: 2ug/mL HRP-Linked Caprine Anti-Rabbit IgG Polyclonal Antibody (Catalog:

**WB (Western Blot)**

Western Blot: Sample: Recombinant IL35, Human.

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/interleukin-35-il35-antibody-western-blot-immunohistochemistry-immunocytochemistry-immunoprecipitation-elisa/20105>