

Rabbit anti-Mouse Monocyte Chemotactic Protein 3 (MCP3) Antibody | anti-MCP3 antibody

Polyclonal Antibody to Monocyte Chemotactic Protein 3 (MCP3)

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA20140	Rabbit	Mouse	Western Blot, Immunohistochemistry, Immunocytochemistry, Immunoprecipitation


PRODUCT SYNONYMS

Monocyte Chemotactic Protein 3 (MCP3); Polyclonal Antibody to Monocyte Chemotactic Protein 3 (MCP3); anti-MCP3 antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Mouse
Purity/Purification	Antigen-specific affinity chromatography followed by Protein A affinity chromatography
Form/Format	Supplied as solution form in 0.01M PBS, pH7.4, containing 0.05% Proclin-300, 50% glycerol.
Concentration	0.24mg/ml (varies by lot)
Applicable Applications for anti-MCP3 antibody	WB (Western Blot), IHC (Immunohistochemistry), ICC (Immunocytochemistry), IP (Immunoprecipitation)
Application Notes	Western blotting: 0.5-2ug/mL Immunohistochemistry: 5-20ug/mL Immunocytochemistry: 5-20ug/mL Optimal working dilutions must be determined by end user.
Source	Polyclonal antibody preparation
Traits	Liquid
Cross Reactivity	Rat
Immunogen	Recombinant MCP3 (Pro28~Pro97) expressed in E.coli .
Preparation and Storage	Storage: Avoid repeated freeze/thaw cycles. Store at 4 degree C for frequent use. Aliquot and store at -20 degree C for 24 months. Stability Test: thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37 degree C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.


IMAGES




IHC (Immunohistochemistry)
DAB staining on IHC-P; Samples: Mouse Testis Tissue.




IHC (Immunohistochemistry)
DAB staining on IHC-P; Samples: Mouse Uterus Tissue.




IHC (Immunohistochemistry)
DAB staining on IHC-P; Samples: Mouse Liver Tissue



IHC (Immunohistochemistry)
DAB staining on IHC-P; Samples: Mouse Ovary Tissue; Primary Ab: 20ug/ml Rabbit Anti-Mouse MCP3 Antibody; Second Ab: 2ug/mL HRP-Linked Caprine Anti-Rabbit IgG Polyclonal Antibody (Catalog:)



IHC (Immunohistochemistry)
DAB staining on IHC-P; Samples: Mouse Stomach Tissue; Primary Ab: 20ug/ml Rabbit Anti-Mouse MCP3 Antibody; Second Ab: 2ug/mL HRP-Linked Caprine Anti-Rabbit IgG Polyclonal Antibody (Catalog:)



WB (Western Blot)
Sample: Recombinant MCP3, Mouse.

NCBI AND UNIPROT INFORMATION

NCBI GI #	7305463
NCBI GeneID	20306
NCBI Accession #	NP_038682.1
NCBI GenBank Nucleotide #	NM_013654.3
UniProt Accession #	Q03366
NCBI Official Full Name	C-C motif chemokine 7

NCBI Official Synonym Full Names	chemokine (C-C motif) ligand 7
NCBI Official Symbol	Ccl7
NCBI Official Synonym Symbols	fic; marc; mcp3; MCP-3; Scya7
NCBI Protein Information	C-C motif chemokine 7
UniProt Protein Name	C-C motif chemokine 7
UniProt Gene Name	Ccl7
UniProt Synonym Gene Names	Fic; Mcp3; Scya7; MCP-3

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/rabbit-mouse-antibody/monocyte-chemotactic-protein-3-mcp3/20140>