

Rabbit CDC25B Polyclonal Antibody | anti-CDC25B antibody

CDC25B antibody - N-terminal region

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA201682	Rabbit	Dog, Guinea Pig, Horse, Pig, Rabbit	Western Blot, Immunohistochemistry


PRODUCT SYNONYMS

CDC25B; Polyclonal Antibody; CDC25B antibody - N-terminal region; anti-CDC25B antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Dog, Guinea Pig, Horse, Pig, Rabbit
Clonality	Polyclonal
Purity/Purification	Affinity Purified
Form/Format	Liquid. Purified antibody supplied in 1x PBS buffer with 0.09% (w/v) sodium azide and 2% sucrose.
Sequence	Synthetic peptide with the following region: ME VPQPEPAPGS ALSPAGVCGG AQRPGHLPGL LLGSHGLLGS PVRAAASS
Sequence Length	580
Applicable Applications for anti-CDC25B antibody	WB (Western Blot), IHC (Immunohistochemistry)
Homology	Dog: 76%; Guinea Pig: 100%; Horse: 90%; Pig: 92%; Rabbit: 84%
Immunogen	The immunogen is a synthetic peptide directed towards the N terminal region of human CDC25B
Preparation and Storage	For short term use, store at 2-8 degree C up to 1 week. For long term storage, store at -20 degree C in small aliquots to prevent freeze-thaw cycles.

IMAGES



WB (Western Blot)

WB Suggested Anti-CDC25B Antibody Titration: 0.2-1 ug/ml Positive Control: Daudi cell lysate CDC25B is supported by BioGPS gene expression data to be expressed in Daudi



IHC (Immunohistochemistry)

Human Liver

ADDITIONAL INFORMATION

Related Product Information for anti-CDC25B antibody	This is a rabbit polyclonal antibody against CDC25B. It was validated on Western Blot and immunohistochemistry Target Description: CDC25B is a member of the CDC25 family of phosphatases. CDC25B activates the cyclin dependent kinase CDC2 by removing two phosphate groups and it is required for entry into mitosis. CDC25B shuttles between the nucleus and the cytoplasm due to nuclear localization and nuclear export signals. The protein is nuclear in the M and G1 phases of the cell cycle and moves to the cytoplasm during S and G2. CDC25B has oncogenic properties, although its role in tumor formation has not been determined.
Product Categories/Family for anti-CDC25B antibody	Polyclonal; Cell Cycle Proteins; Chromatin & Nuclear Signaling; Signal Transduction; Meiosis/Mitosis/Cell Cycle; Immunohistochemistry; Cell Cycle; MAPK Signaling; Phosphorylation

NCBI AND UNIPROT INFORMATION

NCBI GI #	11641413
NCBI GeneID	994
NCBI Accession #	NP_068659
NCBI GenBank Nucleotide #	NM_021873
UniProt Accession #	P30305
Molecular Weight	65kDa
NCBI Official Full Name	M-phase inducer phosphatase 2 isoform 1
NCBI Official Synonym Full Names	cell division cycle 25B
NCBI Official Symbol	CDC25B
NCBI Protein Information	M-phase inducer phosphatase 2
UniProt Protein Name	M-phase inducer phosphatase 2
UniProt Gene Name	CDC25B
UniProt Synonym Gene Names	CDC25HU2
UniProt Entry Name	MPIP2_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/cdc25b-antibody-western-blot-immunohistochemistry-elisa/201682>