

## Human Noggin (NOG) ELISA Kit | NOG elisa kit

Human Noggin (NOG) ELISA Kit

CATALOG #  
**AAA20672**

REACTIVITY  
**Human**


### PRODUCT SYNONYMS

Noggin (NOG); Human Noggin (NOG) ELISA Kit; NOG elisa kit

### PRODUCT SPECIFICATIONS

|                         |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                       |
|-------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Reactivity              | Human                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                 |
| Specificity             | This assay has high sensitivity and excellent specificity for detection of noggin. No significant cross-reactivity or interference between noggin and analogues was observed.                                                                                                                                                                                                                                                                                                                         |
| Sequence Length         | 232                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                   |
| Assay Type              | Quantitative Sandwich                                                                                                                                                                                                                                                                                                                                                                                                                                                                                 |
| Samples                 | Human serum, plasma, tissue homogenates, cell lysates, cell culture supernates and other biological fluids                                                                                                                                                                                                                                                                                                                                                                                            |
| Detection Range         | 15.6-1,000ng/mL                                                                                                                                                                                                                                                                                                                                                                                                                                                                                       |
| Sensitivity             | < 6.0ng/mL                                                                                                                                                                                                                                                                                                                                                                                                                                                                                            |
| Intra-assay Precision   | Intra-assay Precision (Precision within an assay): 3 samples with low, middle and high level noggin were tested 20 times on one plate, respectively. Intra-Assay: CV<10%                                                                                                                                                                                                                                                                                                                              |
| Inter-assay Precision   | Inter-assay Precision (Precision between assays): 3 samples with low, middle and high level noggin were tested on 3 different plates, 8 replicates in each plate. CV(%) = SD/meanX100. Inter-Assay: CV<12%                                                                                                                                                                                                                                                                                            |
| Preparation and Storage | <p>The stability of kit is determined by the loss rate of activity. The loss rate of this kit is less than 5% within the expiration date under appropriate storage condition.</p> <p>To minimize extra influence on the performance, operation procedures and lab conditions, especially room temperature, air humidity, incubator temperature should be strictly controlled. It is also strongly suggested that the whole assay is performed by the same operator from the beginning to the end.</p> |

### IMAGES



Standard Curve (Sample)

ADDITIONAL INFORMATION

|                                               |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                |
|-----------------------------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Related Product Information for NOG elisa kit | <p>Principle of the Assay: The microplate provided in this kit has been pre-coated with an antibody specific to noggin. Standards or samples are then added to the appropriate microplate wells with a biotin-conjugated antibody specific to noggin. Next, Avidin conjugated to Horseradish Peroxidase (HRP) is added to each microplate well and incubated. After TMB substrate solution is added, only those wells that contain noggin, biotin-conjugated antibody and enzyme-conjugated Avidin will exhibit a change in color. The enzyme-substrate reaction is terminated by the addition of sulphuric acid solution and the color change is measured spectrophotometrically at a wavelength of 450nm ± 10nm. The concentration of noggin in the samples is then determined by comparing the O.D. of the samples to the standard curve.</p> <p>Intended Uses: The kit is a sandwich enzyme immunoassay for in vitro quantitative measurement of noggin in human serum, plasma, tissue homogenates, cell lysates, cell culture supernates and other biological fluids.</p> |
| Product Categories/Family for NOG elisa kit   | Signal transduction; Immune molecule; Neuroscience; Bone metabolism                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                            |

NCBI AND UNIPROT INFORMATION

|                                  |                                    |
|----------------------------------|------------------------------------|
| NCBI GI #                        | 4885523                            |
| NCBI GeneID                      | 9241                               |
| NCBI Accession #                 | NP_005441.1                        |
| NCBI GenBank Nucleotide #        | NM_005450.4                        |
| UniProt Accession #              | Q13253                             |
| Molecular Weight                 | 25,774 Da                          |
| NCBI Official Full Name          | noggin                             |
| NCBI Official Synonym Full Names | noggin                             |
| NCBI Official Symbol             | NOG                                |
| NCBI Official Synonym Symbols    | SYM1; SYNS1                        |
| NCBI Protein Information         | noggin; symphalangism 1 (proximal) |
| UniProt Protein Name             | Noggin                             |
| UniProt Gene Name                | NOG                                |
| UniProt Entry Name               | NOGG_HUMAN                         |

**For Research Use Only.** Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.