

Rabbit anti-Human Syndecan 1 Polyclonal Antibody | anti-SDC1 antibody

Polyclonal Antibody to Syndecan 1 (SDC1)

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA20851	Rabbit	Human	Western Blot, Immunohistochemistry, Immunocytochemistry, Immunoprecipitation


PRODUCT SYNONYMS

Syndecan 1; Polyclonal Antibody; Polyclonal Antibody to Syndecan 1 (SDC1); anti-SDC1 antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Purity/Purification	Antigen-specific affinity chromatography followed by Protein A affinity chromatography
Form/Format	Supplied as solution form in PBS, pH7.4, containing 0.02% NaN ₃ , 50% glycerol.
Concentration	1mg/ml (varies by lot)
Applicable Applications for anti-SDC1 antibody	WB (Western Blot), IHC (Immunohistochemistry), ICC (Immunocytochemistry), IP (Immunoprecipitation)
Application Notes	Western blotting: 0.5-2ug/mL Immunohistochemistry: 5-20ug/mL Immunocytochemistry: 5-20ug/mL Optimal working dilutions must be determined by end user.
Source	Polyclonal antibody preparation
Traits	Liquid
Immunogen	Recombinant SDC1 (Gln23~Leu253) expressed in E.coli
Preparation and Storage	Storage: Avoid repeated freeze/thaw cycles. Store at 4 degree C for frequent use. Aliquot and store at -20 degree C for 24 months. Stability Test: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37 degree C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.


IMAGES




IHC (Immunohistochemistry)
DAB staining on IHC-P; Samples: Human Liver Tissue.




IHC (Immunohistochemistry)
DAB staining on IHC-P; Samples: Human Kidney Tissue.




IHC (Immunohistochemistry)
DAB staining on IHC-P; Samples: Human Lung Cancer Tissue.



IHC (Immunohistochemistry)
DAB staining on IHC-P; Samples: Human Colorectal cancer Tissue; Primary Ab: 20ug/ml Rabbit Anti-Human SDC1 Antibody; Second Ab: 2ug/mL HRP-Linked Caprine Anti-Rabbit IgG Polyclonal Antibody



IHC (Immunohistochemistry)
DAB staining on IHC-P; Samples: Human Pancreas Tissue; Primary Ab: 20ug/ml Rabbit Anti-Human SDC1 Antibody; Second Ab: 2ug/mL HRP-Linked Caprine Anti-Rabbit IgG Polyclonal Antibody



WB (Western Blot)
Western Blot: Sample: Recombinant SDC1, Human.

NCBI AND UNIPROT INFORMATION

NCBI GI #	55749480
NCBI GeneID	6382
NCBI Accession #	NP_001006947.1
NCBI GenBank Nucleotide #	NM_001006946.1
UniProt Accession #	P18827; Q53QV0; Q546D3; Q96HB7; D6W523; AAB21434; AAB23935
NCBI Official Full Name	syndecan-1

NCBI Official Synonym Full Names	syndecan 1
NCBI Official Symbol	SDC1
NCBI Official Synonym Symbols	SDC; CD138; SYND1; syndecan
NCBI Protein Information	syndecan-1
UniProt Protein Name	Syndecan-1
UniProt Gene Name	SDC1
UniProt Synonym Gene Names	SDC; SYND1

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/syndecan-1-antibody-western-blot-immunohistochemistry-immunocytochemistry-immunoprecipitation-elisa/20851>