

Rabbit anti-Human BHC110/LSD1 Polyclonal Antibody | anti-BHC110/LSD1 antibody

Rabbit anti-BHC110/LSD1 Antibody, Affinity Purified

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA210822	Rabbit	Human	Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

BHC110/LSD1; Polyclonal Antibody; Rabbit anti-BHC110/LSD1 Antibody; Affinity Purified; flavin-containing amine oxidase domain-containing protein 2; lysine-specific histone demethylase 1; lysine-specific histone demethylase 1A; lysine (K)-specific demethylase 1A; LSD1; KDM1; FAD-binding protein BRAF35-HDAC complex; 110 kDa subunit; CPRF; BRAF35-HDAC complex protein BHC110; BHC110; amine oxidase (flavin containing) domain 2; AOF2; Lysine-specific histone demethylase 1A; lysine demethylase 1A; anti-BHC110/LSD1 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Purity/Purification	Antigen Affinity Purified
Form/Format	Liquid. Whole IgG. Tris-citrate/phosphate buffer, pH 7 to 8 containing 0.09% Sodium Azide
Concentration	1000ug/ml (varies by lot)
Applicable Applications for anti-BHC110/LSD1 antibody	IHC (Immunohistochemistry), WB (Western Blot)
Predicted Reactivity	Mouse
Immunogen	Between 802 and C-terminus
Conjugation	Unconjugated
Preparation and Storage	Store at 2-8 degree C Shelf Life: 1 year from date of receipt

IMAGES



WB (Western Blot)

Detection of human BHC110/LSD1 by western blot. Samples: Whole cell lysate (50 ug) from HeLa and HEK293T cells prepared using NETN lysis buffer. Antibody: Affinity purified rabbit anti-BHC110/LSD1 antibody AAA210822 (lot AAA210822-1) used for WB at 0.06 ug/ml. Detection: Chemiluminescence with an exposure time of 3 minutes.



IHC (Immunohistochemistry)

Detection of human BHC110/LSD1 by immunohistochemistry. Sample: FFPE section of human breast carcinoma. Antibody: Affinity purified rabbit anti-BHC110/LSD1 (Cat. No. AAA210822) used at a dilution of 1:1,000 (1ug/ml). Detection: DAB

ADDITIONAL INFORMATION

Related Product Information for anti-BHC110/LSD1 antibody	BHC110/LSD1 is a histone demethylase and a component of several histone deacetylase complexes. BHC110/LSD1 is able to specifically catalyze the demethylation of Lys4 of histone H3. As part of histone deacetylase complexes BHC110/LSD1 functions as a corepressor of transcription. BHC110/LSD1 has also been demonstrated to interact with the androgen receptor and stimulate androgen-receptor-dependent transcription.
Product Categories/Family for anti-BHC110/LSD1 antibody	Primary Antibodies

NCBI AND UNIPROT INFORMATION

NCBI GI #	226437582
NCBI GeneID	23028
NCBI Accession #	NP_055828.2
NCBI GenBank Nucleotide #	NM_015013
UniProt Accession #	O60341; Q5TH94; Q5TH95; Q86VT7; Q8IXK4; Q8NDP6; Q8TAZ3; Q96AW4; A8MWP9
Molecular Weight	95,155 Da
NCBI Official Full Name	Homo sapiens lysine (K)-specific demethylase 1A (KDM1A), transcript variant 2, mRNA
NCBI Official Synonym Full Names	lysine (K)-specific demethylase 1A
NCBI Official Symbol	KDM1A
NCBI Official Synonym Symbols	AOF2; KDM1; LSD1; BHC110
NCBI Protein Information	lysine-specific histone demethylase 1A; BRAF35-HDAC complex protein BHC110; FAD-binding protein BRAF35-HDAC complex, 110 kDa subunit; amine oxidase (flavin containing) domain 2; flavin-containing amine oxidase domain-containing protein 2; lysine (K)-speci
UniProt Protein Name	Lysine-specific histone demethylase 1A
UniProt Gene Name	KDM1A
UniProt Synonym Gene Names	AOF2; KDM1; KIAA0601; LSD1
UniProt Entry Name	KDM1A_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/bhc110-lsd1-antibody-immunohistochemistry-western-blot-elisa/210822>