

Rabbit anti-Human, Mouse WDR20 Polyclonal Antibody | anti-WDR20 antibody

Rabbit anti-WDR20 Antibody, Affinity Purified

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA211573	Rabbit	Human, Mouse	Immunohistochemistry, Immunoprecipitation, Western Blot


PRODUCT SYNONYMS

WDR20; Polyclonal Antibody; Rabbit anti-WDR20 Antibody; Affinity Purified; DMR; Protein DMR; WD repeat-containing protein 20; WD repeat domain 20; anti-WDR20 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human, Mouse
Clonality	Polyclonal
Purity/Purification	Antigen Affinity Purified
Form/Format	Liquid. Whole IgG. Tris-citrate/phosphate buffer, pH 7 to 8 containing 0.09% Sodium Azide
Concentration	1000ug/ml (varies by lot)
Applicable Applications for anti-WDR20 antibody	IHC (Immunohistochemistry), IP (Immunoprecipitation), WB (Western Blot)
Immunogen	Between 450 and 500
Conjugation	Unconjugated
Preparation and Storage	Store at 2-8 degree C Shelf Life: 1 year from date of receipt

IMAGES



WB (Western Blot)

Detection of human and mouse WDR20 by western blot (h&m) and immunoprecipitation (h).
Samples: Whole cell lysate from HeLa (5, 15 and 50 ug for WB; 1 mg for IP, 20% of IP loaded), HEK293T (T; 50 ug), and mouse NIH 3T3 (M; 50 ug) cells. Antibodies: Affinity purified rabbit anti-WDR20 antibody AAA211573 used for WB at 0.1 ug/ml (A) and 1 ug/ml (B) and used for IP at 3 ug/mg lysate. WDR20 was also immunoprecipitated by rabbit anti-WDR20 antibody BL6027, which recognizes an upstream epitope. Detection: Chemiluminescence with exposure times of 30 seconds (A) and 10 seconds (B).



IHC (Immunohistochemistry)

Detection of human WDR20 by immunohistochemistry. Sample: FFPE section of human small cell lung cancer. Antibody: Affinity purified rabbit anti-WDR20 (Cat. No. AAA211573 Lot1) used at a dilution of 1:1,000 (1ug/ml). Detection: DAB

ADDITIONAL INFORMATION

Related Product Information for anti-WDR20 antibody	WDR20 (WD repeat-containing protein 20) contains five WD repeats and is a member of the WD family of proteins. The WD repeat is defined by four or more repeating units of a conserved core of approximately 40 amino acids ending with tryptophan-aspartic acid (WD). WD repeats may serve as sites of protein-protein interaction for adaptor proteins and facilitate multiprotein complex formation. WD proteins are involved in a variety of cellular processes. The function of WDR20 has not been characterized.
---	--

NCBI AND UNIPROT INFORMATION

NCBI GeneID	91833
UniProt Accession #	Q8TBZ3
UniProt Protein Name	WD repeat-containing protein 20
UniProt Gene Name	WDR20
UniProt Entry Name	WDR20_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/polyclonal/wdr20-antibody-immunohistochemistry-immunoprecipitation-western-blot-elisa/211573>