

Rabbit anti-Human Cyclin T1 Polyclonal Antibody | anti-CCNT1 antibody

Rabbit anti-Cyclin T1 Antibody, Affinity Purified

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA212191	Rabbit	Human	Immunohistochemistry, Immunoprecipitation, Western Blot

PRODUCT SYNONYMS

Cyclin T1; Polyclonal Antibody; Rabbit anti-Cyclin T1 Antibody; Affinity Purified; CYCT1; cyclin-T1; HIVE1; MLLT10/CCNT1 fusion; cyclin C-related protein; CDK9-associated C-type protein; CCNT; human immunodeficiency virus type 1 (HIV-1) expression (elevated) 1; Cyclin-T; Cyclin-T1; cyclin T1; anti-CCNT1 antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Purity/Purification	Antigen Affinity Purified
Form/Format	Liquid. Whole IgG. Tris-buffered Saline containing 0.1% BSA and 0.09% Sodium Azide
Concentration	200ug/ml (varies by lot)
Applicable Applications for anti-CCNT1 antibody	IHC (Immunohistochemistry), IP (Immunoprecipitation), WB (Western Blot)
Predicted Reactivity	Bovine, Horse, Chimpanzee
Immunogen	Between 425 and 475
Conjugation	Unconjugated
Preparation and Storage	Store at 2-8 degree C Shelf Life: 1 year from date of receipt

IMAGES




WB (Western Blot)

Detection of human Cyclin T1 by western blot. Samples: Whole cell lysate from HeLa (15 and 50 ug) and HEK293T (T; 50 ug) cells. Antibody: Affinity purified rabbit anti-Cyclin T1 antibody AAA212191 (lot AAA212191-1) used for IP at 6 ug/mg lysate. Cyclin T1 was also used at 0.04 ug/ml. Detection: Chemiluminescence with an exposure time of 30 seconds.



IP (Immunoprecipitation)

Detection of human Cyclin T1 by western blot of immunoprecipitates. Samples: Whole cell lysate (1 mg for IP, 20% of IP loaded) from HeLa cells. Antibodies: Affinity purified rabbit anti-Cyclin T1 antibody AAA212191 (lot AAA212191-1) used for IP at 6 ug/mg lysate. Cyclin T1 was also immunoprecipitated by rabbit anti-Cyclin T1 antibodies



IHC (Immunohistochemistry)

Detection of human Cyclin T1 by immunohistochemistry. Sample: FFPE section of human breast carcinoma. Antibody: Affinity purified rabbit anti-Cyclin T1 (Cat. No. AAA212191 lot 1) used at a dilution of 1:200 (1ug/ml). Detection: DAB

ADDITIONAL INFORMATION

Related Product Information for anti-CCNT1 antibody	Cyclin T1 is a regulatory subunit of the cyclin-dependent kinase pair (CDK9/cyclin-T1) complex, also called positive transcription elongation factor B (P-TEFb), which is proposed to facilitate the transition from abortive to productive elongation by phosphorylating the CTD (carboxy-terminal domain) of the large subunit of RNA polymerase II (RNA Pol II). In case of HIV or SIV infections, Cyclin T1 binds to the transactivation domain of the viral nuclear transcriptional activator, Tat, thereby increasing Tat's affinity for the transactivating response RNA element (TAR RNA) [taken from the Universal Protein Resource (UniProt) www.uniprot.org/uniprot/O60563].
Product Categories/Family for anti-CCNT1 antibody	Primary Antibodies

NCBI AND UNIPROT INFORMATION

NCBI GI #	17978466
NCBI GeneID	904
NCBI Accession #	NP_001231.2
NCBI GenBank Nucleotide #	NM_001240.3
UniProt Accession #	O60563 ; O60581 ; A9XU13 ; E7EX76
Molecular Weight	21,206 Da
NCBI Official Full Name	cyclin-T1 isoform a
NCBI Official Synonym Full Names	cyclin T1
NCBI Official Symbol	CCNT1
NCBI Official Synonym Symbols	CCNT; CYCT1; HIVE1
NCBI Protein Information	cyclin-T1
UniProt Protein Name	Cyclin-T1
UniProt Gene Name	CCNT1
UniProt Synonym Gene Names	CycT1; Cyclin-T
UniProt Entry Name	CCNT1_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/cyclin-t1-antibody-immunohistochemistry-immunoprecipitation-western-blot-elisa/212191>