

Goat anti-Human, Mouse H2AX Polyclonal Antibody | anti-H2AX antibody

Goat anti-H2AX Antibody, Affinity Purified

CATALOG # AAA212305	HOST Goat	REACTIVITY Human, Mouse	APPLICATIONS Immunohistochemistry, Western Blot
-------------------------------	---------------------	-----------------------------------	---

PRODUCT SYNONYMS

H2AX; Polyclonal Antibody; Goat anti-H2AX Antibody; Affinity Purified; H2AX histone; histone H2AX; H2A/X; H2A.X; histone H2A.x; Histone H2AX; H2A histone family member X; anti-H2AX antibody

PRODUCT SPECIFICATIONS

Host	Goat
Reactivity	Human, Mouse
Clonality	Polyclonal
Purity/Purification	Antigen Affinity Purified
Form/Format	Liquid. Whole IgG. Tris-citrate/phosphate buffer, pH 7 to 8 containing 0.09% Sodium Azide
Concentration	1000ug/ml (varies by lot)
Applicable Applications for anti-H2AX antibody	IHC (Immunohistochemistry), WB (Western Blot)
Immunogen	Between 93 and C-terminus
Conjugation	Unconjugated
Preparation and Storage	Store at 2-8 degree C Shelf Life: 1 year from date of receipt

IMAGES




WB (Western Blot)

Detection of human H2AX by western blot. Samples: Whole cell lysate (50 ug) from HeLa, HEK293T, Jurkat, and K-562 cells prepared using RIPA lysis buffer. Antibody: Affinity purified goat anti-H2AX antibody AAA212305 (lot AAA212305-2) used for WB at 0.1 ug/ml. Detection: Chemiluminescence with an exposure time of 30 seconds.



IHC (Immunohistochemistry)

Detection of human H2AX by immunohistochemistry. Sample: FFPE section of human ovarian cancer. Antibody: Affinity purified goat anti-H2AX (Cat. No. AAA212305 lot 2) used at a dilution of 1:5,000 (0.2ug/ml). Detection: DAB



IHC (Immunohistochemistry)

Detection of mouse H2AX by immunohistochemistry. Sample: FFPE section of mouse CT26 colon carcinoma. Antibody: Affinity purified goat anti-H2AX (Cat. No. AAA212305 lot 2) used at a dilution of 1:1,000 (1ug/ml). Detection: DAB

ADDITIONAL INFORMATION

Related Product Information for anti-H2AX antibody	H2AX is a member of the histone H2A family. The four core histones involved in the formation of the nucleosome structure of compacted chromatin are H2A, H2B, H3, and H4. H2AX may function to facilitate DNA repair, and recent studies have shown that H2AX is required for the maintenance of genomic stability.
Product Categories/Family for anti-H2AX antibody	Primary Antibodies

NCBI AND UNIPROT INFORMATION

NCBI GI #	4504253
NCBI GeneID	3014
NCBI Accession #	NP_002096.1
NCBI GenBank Nucleotide #	NM_002105.2
UniProt Accession #	P16104; Q4ZGJ7; Q6IAS5
Molecular Weight	15,145 Da
NCBI Official Full Name	histone H2A.x
NCBI Official Synonym Full Names	H2A histone family, member X
NCBI Official Symbol	H2AFX
NCBI Official Synonym Symbols	H2AX; H2A.X; H2A/X
NCBI Protein Information	histone H2A.x; H2AX histone
UniProt Protein Name	Histone H2A.x
UniProt Gene Name	H2AFX
UniProt Synonym Gene Names	H2AX; H2a/x
UniProt Entry Name	H2AX_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/h2ax-antibody-immunohistochemistry-western-blot-elisa/212305>