

Rabbit anti-Human MKI67IP Polyclonal Antibody | anti-MKI67IP antibody

Rabbit anti-MKI67IP Antibody, Affinity Purified

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA212475	Rabbit	Human	Immunoprecipitation, Western Blot


PRODUCT SYNONYMS

MKI67IP; Polyclonal Antibody; Rabbit anti-MKI67IP Antibody; Affinity Purified; nucleolar phosphoprotein Nopp34; nucleolar protein interacting with the FHA domain of pKi-67; Nopp34; MKI67 FHA domain-interacting nucleolar phosphoprotein; hNIFK; nucleolar protein interacting with the FHA domain of MKI67; anti-MKI67IP antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Purity/Purification	Antigen Affinity Purified
Form/Format	Liquid. Whole IgG. Tris-citrate/phosphate buffer, pH 7 to 8 containing 0.09% Sodium Azide
Concentration	1000ug/ml (varies by lot)
Applicable Applications for anti-MKI67IP antibody	IP (Immunoprecipitation), WB (Western Blot)
Immunogen	Between 243 and C-terminus
Conjugation	Unconjugated
Preparation and Storage	Store at 2-8 degree C Shelf Life: 1 year from date of receipt

IMAGES



WB (Western Blot)

Detection of human MKI67IP by western blot. Samples: Whole cell lysate (50 ug) from HEK293T, HeLa, and Jurkat cells. Antibodies: Affinity purified rabbit anti-MKI67IP antibody AAA212475 (lot AAA212475-1) used for WB at 0.04 ug/ml. Detection: Chemiluminescence with an exposure time of 10 seconds.



IP (Immunoprecipitation)

Detection of human MKI67IP by western blot of immunoprecipitates. Samples: Whole cell lysate (1 mg for IP; 20% of IP loaded) from HEK293T cells. Antibodies: Affinity purified rabbit anti-MKI67IP antibody AAA212475 (lot AAA212475-1) used for IP at 6 ug/mg lysate. MKI67IP was also immunoprecipitated by rabbit anti-MKI67IP antibody BL14642. For blotting immunoprecipitated MKI67IP, AAA212475 was used at 0.2 ug/ml. Detection: Chemiluminescence with an exposure time of 3 seconds.

ADDITIONAL INFORMATION

Related Product Information for anti-MKi67IP antibody	MKi67 (FHA domain) interacting nucleolar phosphoprotein (MKi67IP) is a protein that interacts with the forkhead-associated domain of the Ki-67 antigen. The encoded protein may bind RNA and may play a role in mitosis and cell cycle progression [taken from NCBI Entrez Gene (Gene ID: 84365)].
Product Categories/Family for anti-MKi67IP antibody	Primary Antibodies

NCBI AND UNIPROT INFORMATION

NCBI GI #	222352111
NCBI GeneID	84365
NCBI Accession #	NP_115766.3
NCBI GenBank Nucleotide #	NM_032390.4
UniProt Accession #	Q9BYG3; Q8TB66; Q96ED4; A8K788
Molecular Weight	34,222 Da
NCBI Official Full Name	MKi67 FHA domain-interacting nucleolar phosphoprotein
NCBI Official Synonym Full Names	MKi67 (FHA domain) interacting nucleolar phosphoprotein
NCBI Official Symbol	MKi67IP
NCBI Official Synonym Symbols	NIFK; Nopp34
NCBI Protein Information	MKi67 FHA domain-interacting nucleolar phosphoprotein; hNIFK; nucleolar phosphoprotein Nopp34; nucleolar protein interacting with the FHA domain of pKi-67
UniProt Protein Name	MKi67 FHA domain-interacting nucleolar phosphoprotein
UniProt Gene Name	MKi67IP
UniProt Synonym Gene Names	NIFK; NOPP34; hNIFK
UniProt Entry Name	MK67I_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/mki67ip-antibody-immunoprecipitation-western-blot-elisa/212475>