

Rabbit anti-Human, Mouse FAF2/ETEA Polyclonal Antibody | anti-FAF2/ETEA antibody

Rabbit anti-FAF2/ETEA Antibody, Affinity Purified


CATALOG # AAA213238	HOST Rabbit	REACTIVITY Human, Mouse	APPLICATIONS Immunoprecipitation, Western Blot
-------------------------------	-----------------------	-----------------------------------	--

PRODUCT SYNONYMS

FAF2/ETEA; Polyclonal Antibody; Rabbit anti-FAF2/ETEA Antibody; Affinity Purified; Protein ETEA; UBXD8; UBX domain-containing protein 8; UBX domain-containing protein 3B; UBXN3B; FAS-associated factor 2; expressed in T-cells and eosinophils in atopic dermatitis; ETEA; UBX domain protein 3B; UBX domain containing 8; Fas associated factor family member 2; anti-FAF2/ETEA antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse
Clonality	Polyclonal
Purity/Purification	Antigen Affinity Purified
Form/Format	Liquid. Whole IgG. Tris-citrate/phosphate buffer, pH 7 to 8 containing 0.09% Sodium Azide
Concentration	1000ug/ml (varies by lot)
Applicable Applications for anti-FAF2/ETEA antibody	IP (Immunoprecipitation), WB (Western Blot)
Predicted Reactivity	Rat, Bovine
Immunogen	Between 395 and C-terminus
Conjugation	Unconjugated
Preparation and Storage	Store at 2-8 degree C Shelf Life: 1 year from date of receipt



WB (Western Blot)

Detection of human and mouse FAF2 by western blot. Samples: Whole cell lysate (50 ug) from HeLa, HEK293T, Jurkat, mouse TCMK-1, and mouse NIH 3T3 cells prepared using NETN lysis buffer. Antibody: Affinity purified rabbit anti-FAF2 antibody AAA213238 (lot AAA213238-1) used for WB at 0.1 ug/ml. Detection: Chemiluminescence with an exposure time of 30 seconds.



IP (Immunoprecipitation)

Detection of human FAF2 by western blot of immunoprecipitates. Samples: Whole cell lysate (0.5 or 1.0 mg per IP reaction; 20% of IP loaded) from 293T cells prepared using NETN lysis buffer. Antibodies: Affinity purified rabbit anti-FAF2 antibody AAA213238 (lot AAA213238-1) used for IP at 6 ug per reaction. FAF2 was also immunoprecipitated by rabbit anti-FAF2 antibody For blotting immunoprecipitated FAF2, AAA213238 was used at 0.4 ug/ml. Detection: Chemiluminescence with an exposure time of 10 seconds.

ADDITIONAL INFORMATION

Related Product Information for anti-FAF2/ETEA antibody	FAF2/ETEA is highly expressed in peripheral blood of patients with atopic dermatitis (AD), compared to normal individuals. It may play a role in regulating the resistance to apoptosis that is observed in T cells and eosinophils of AD patients [taken from NCBI Entrez Gene (Gene ID: 23197)].
Product Categories/Family for anti-FAF2/ETEA antibody	Primary Antibodies

NCBI AND UNIPROT INFORMATION

NCBI GI #	24797106
NCBI GeneID	23197
NCBI Accession #	NP_055428.1
NCBI GenBank Nucleotide #	NM_014613.2
UniProt Accession #	Q96CS3
NCBI Official Full Name	FAS-associated factor 2
NCBI Official Synonym Full Names	Fas associated factor family member 2
NCBI Official Symbol	FAF2
NCBI Official Synonym Symbols	ETEA; UBXD8; UBXN3B
NCBI Protein Information	FAS-associated factor 2
UniProt Protein Name	FAS-associated factor 2
UniProt Gene Name	FAF2
UniProt Synonym Gene Names	ETEA; KIAA0887; UBXD8; UBXN3B
UniProt Entry Name	FAF2_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

