

IgG Heavy and Light Chain Cross-Adsorbed Secondary Antibody | IgG secondary antibody

Rabbit anti-Mouse IgG Heavy and Light Chain Cross-Adsorbed Antibody DyLight 488 Conjugated

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA213715	Rabbit	Mouse Minimum Reactivity: Human, Rat	Immunofluorescence, Flow Cytometry, Functional Assay, Immunocytochemistry, Immunohistochemistry, Western Blot


PRODUCT SYNONYMS

IgG Heavy and Light Chain Cross-Adsorbed; Polyclonal Secondary Antibody; Rabbit anti-Mouse IgG Heavy and Light Chain Cross-Adsorbed Antibody DyLight 488 Conjugated; Mouse IgG-Heavy and Light Chain; IgG secondary antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Mouse Minimum Reactivity: Human, Rat
Clonality	Polyclonal
Purity/Purification	Antigen Affinity Purified
Form/Format	Liquid. Whole IgG. Phosphate Buffered Saline (PBS) containing 0.2% BSA and 0.09% Sodium Azide
Concentration	0.5mg/ml (varies by lot)
Applicable Applications for IgG secondary antibody	IF (Immunofluorescence), FCM/FACS (Flow Cytometry), ICC (Immunocytochemistry), IHC (Immunohistochemistry), WB (Western Blot)
Conjugation	DyLight 488
Preparation and Storage	Store at 2-8 degree C Shelf Life: 1 year from date of receipt

IMAGES



WB (Western Blot)

Detection of GAPDH and hnRNP in HeLa Whole Cell Lysate. Primary Antibodies: cocktail of goat anti-GAPDH and mouse anti-hnRNP at 1 ug/ml each. Secondary Antibodies: cocktail of DyLight 680-conjugated rabbit anti-goat (red) and DyLight 488-conjugated rabbit anti-mouse AAA213715 (AAA213715-3) (blue) at 0.5 ug/ml each. Acquisition: Syngene G:Box, 39 seconds (red) and 60 seconds (blue).



IF (Immunofluorescence)

Detection of human p53 by immunofluorescence. Sample: FFPE section of human breast carcinoma. Primary Antibody: mouse anti-p53 (clone DO-1) used at a dilution of 1:100. Secondary Antibody: Green-fluorescent Rabbit anti-mouse IgG-heavy and light chain cross-adsorbed Antibody DyLight 488 Conjugated (AAA213715 Lot 6) used at a dilution of 1:100 (5ug/ml). Counterstain: DAPI (blue)

ADDITIONAL INFORMATION

Related Product Information for IgG secondary antibody	Anti-heavy- and light-chain antibodies are designed to react with the whole intact Ig molecule. Found in all body fluids and a primary means of protection against infection, an anti-H+L IgG antibody allows for some potential cross-reactivity to other Ig molecules and IgG molecules from other closely-related species.
Product Categories/Family for IgG secondary antibody	Secondary Antibody Conjugates ; Secondary Antibodies

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/igg-heavy-and-light-chain-cross-adsorbed-secondary-antibody-immunofluorescence-flow-cytometry-immunocytochemistry-immunohistochemistry-western-blot-elisa/213715>