

Rabbit anti-Human BHC110/LSD1 Polyclonal Antibody | anti-BHC110/LSD1 antibody

Rabbit anti-BHC110/LSD1 IHC Antibody, Affinity Purified

CATALOG # AAA213807	HOST Rabbit	REACTIVITY Human	APPLICATIONS Immunohistochemistry
-------------------------------	-----------------------	----------------------------	---


PRODUCT SYNONYMS

BHC110/LSD1; Polyclonal Antibody; Rabbit anti-BHC110/LSD1 IHC Antibody; Affinity Purified; flavin-containing amine oxidase domain-containing protein 2; lysine-specific histone demethylase 1; lysine-specific histone demethylase 1A; lysine (K)-specific demethylase 1A; LSD1; KDM1; FAD-binding protein BRAF35-HDAC complex; 110 kDa subunit; CPRF; BRAF35-HDAC complex protein BHC110; BHC110; amine oxidase (flavin containing) domain 2; AOF2; Lysine-specific histone demethylase 1A; lysine demethylase 1A; anti-BHC110/LSD1 antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Purity/Purification	Antigen Affinity Purified
Form/Format	Liquid. Whole IgG. Tris-buffered Saline containing 0.1% BSA and 0.09% Sodium Azide
Concentration	250ug/ml (varies by lot)
Applicable Applications for anti-BHC110/LSD1 antibody	IHC (Immunohistochemistry)
Predicted Reactivity	Orangutan, Gorilla
Immunogen	Between 25 and 75
Conjugation	Unconjugated
Preparation and Storage	Store at 2-8 degree C Shelf Life: 1 year from date of receipt

IMAGES



IHC (Immunohistochemistry)

Detection of human BHC110/LSD1 by immunohistochemistry. Sample: FFPE section of human breast adenocarcinoma. Antibody: Affinity purified rabbit anti-BHC110/LSD1 (Cat. No. AAA213807) used at a dilution of 1:250. Detection: DAB.



IHC (Immunohistochemistry)

Detection of human BHC110/LSD1 by immunohistochemistry. Sample: FFPE section of human skin basal cell carcinoma. Antibody: Affinity purified rabbit anti-BHC110/LSD1 (Cat. No. AAA213807) used at a dilution of 1:250. Detection: DAB

ADDITIONAL INFORMATION

Related Product Information for anti-BHC110/LSD1 antibody	BHC110/LSD1 is a histone demethylase and a component of several histone deacetylase complexes. BHC110/LSD1 is able to specifically catalyze the demethylation of Lys4 of histone H3. As part of histone deacetylase complexes BHC110/LSD1 functions as a corepressor of transcription. BHC110/LSD1 has also been demonstrated to interact with the androgen receptor and stimulate androgen-receptor-dependent transcription.
Product Categories/Family for anti-BHC110/LSD1 antibody	Primary Antibodies

NCBI AND UNIPROT INFORMATION

NCBI GI #	226437582
NCBI GeneID	23028
NCBI Accession #	NP_055828.2
NCBI GenBank Nucleotide #	NM_015013
UniProt Accession #	O60341; Q5TH94; Q5TH95; Q86VT7; Q8IXK4; Q8NDP6; Q8TAZ3; Q96AW4; A8MWP9
Molecular Weight	95,155 Da
NCBI Official Full Name	Homo sapiens lysine (K)-specific demethylase 1A (KDM1A), transcript variant 2, mRNA
NCBI Official Synonym Full Names	lysine (K)-specific demethylase 1A
NCBI Official Symbol	KDM1A
NCBI Official Synonym Symbols	AOF2; KDM1; LSD1; BHC110
NCBI Protein Information	lysine-specific histone demethylase 1A; BRAF35-HDAC complex protein BHC110; FAD-binding protein BRAF35-HDAC complex, 110 kDa subunit; amine oxidase (flavin containing) domain 2; flavin-containing amine oxidase domain-containing protein 2; lysine (K)-speci
UniProt Protein Name	Lysine-specific histone demethylase 1A
UniProt Gene Name	KDM1A
UniProt Synonym Gene Names	AOF2; KDM1; KIAA0601; LSD1
UniProt Entry Name	KDM1A_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/bhc110-lsd1-antibody-immunohistochemistry-elisa/213807>