

Rabbit anti-Human Cyclin C Polyclonal Antibody | anti-CCNC antibody

Rabbit anti-Cyclin C IHC Antibody, Affinity Purified

CATALOG # AAA214094	HOST Rabbit	REACTIVITY Human	APPLICATIONS Immunohistochemistry
-------------------------------	-----------------------	----------------------------	---


PRODUCT SYNONYMS

Cyclin C; Polyclonal Antibody; Rabbit anti-Cyclin C IHC Antibody; Affinity Purified; SRB11 homolog; SRB11; CycC; hSRB11; cyclin-C; Cyclin-C; cyclin C; anti-CCNC antibody

PRODUCT SPECIFICATIONS


Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Purity/Purification	Antigen Affinity Purified
Form/Format	Liquid. Whole IgG. Tris-buffered Saline containing 0.1% BSA and 0.09% Sodium Azide
Concentration	100ug/ml (varies by lot)
Applicable Applications for anti-CCNC antibody	IHC (Immunohistochemistry)
Predicted Reactivity	Mouse, Rat, Chicken, Turkey, Bovine, Dog, Horse, Rabbit, Pig, Panda, Orangutan, Monkey, Gorilla, Chimpanzee, White-tufted-ear marmoset, Duckbill platypus, African elephant, Small-eared galago, Northern white-cheeked gibbon, Naked mole rat, Thirteen-lined ground squirrel, Chinese hamster, Crab-eating macaque, Tasmanian devil, Gray short-tailed opossum, Little brown bat, Zebra finch
Immunogen	Between 233 and C-terminus
Conjugation	Unconjugated
Preparation and Storage	Store at 2-8 degree C Shelf Life: 1 year from date of receipt

IMAGES



IHC (Immunohistochemistry)

Detection of human Cyclin C by immunohistochemistry. Sample: FFPE section of human pancreatic islet cell tumor. Antibody: Affinity purified rabbit anti-Cyclin C (Cat. No. AAA214094) used at a dilution of 1:100. Detection: DAB



IHC (Immunohistochemistry)

Detection of human Cyclin C by immunohistochemistry. Sample: FFPE section of human skin squamous cell carcinoma. Antibody: Affinity purified rabbit anti-Cyclin C (Cat. No. AAA214094) used at a dilution of 1:100. Detection: DAB

ADDITIONAL INFORMATION

Related Product Information for anti-CCNC antibody	Cyclin C is a member of the cyclin family of regulatory proteins that associate with and activate cyclin-dependent kinases (CDKs) to regulate cell cycle progression. Cyclins exhibit periodicity in their expression and activity which is regulated transcriptionally and post-transcriptionally via phosphorylation and ubiquitination. Cyclin C associates with cyclin-dependent kinase 8 (CDK8). It is also a component of the Mediator complex, a coactivator complex that provides a scaffold to link gene-specific transcription factors to RNA polymerase II and the general transcription machinery.
Product Categories/Family for anti-CCNC antibody	Primary Antibodies

NCBI AND UNIPROT INFORMATION

NCBI GI #	61676090
NCBI GeneID	892
NCBI Accession #	NP_005181.2
NCBI GenBank Nucleotide #	NM_005190
UniProt Accession #	P24863; Q9H543; B4DPZ1
Molecular Weight	~55 kDa
NCBI Official Full Name	Homo sapiens cyclin C (CCNC), transcript variant 1, mRNA
NCBI Official Synonym Full Names	cyclin C
NCBI Official Symbol	CCNC
NCBI Official Synonym Symbols	CycC
NCBI Protein Information	cyclin-C; hSRB11; SRB11 homolog
UniProt Protein Name	Cyclin-C
UniProt Gene Name	CCNC
UniProt Synonym Gene Names	hSRB11
UniProt Entry Name	CCNC_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/polyclonal/cyclin-c-antibody-immunohistochemistry-elisa/214094>