

IL3 active protein

Recombinant Human IL3

CATALOG #
AAA214236

HOST
E Coli


PRODUCT SYNONYMS

IL3; Recombinant Human IL3; IL-3; Mast cell growth factor; MCGF; Multi-CSF; Hematopoietic growth factor; HCGF; P-cell stimulation factor; Recombinant Human Interleukin-3 (IL3); IL3 active protein


PRODUCT SPECIFICATIONS

Host	E Coli
Purity/Purification	>95% as determined by SDS-PAGE
Form/Format	Lyophilized from a 0.2 um filtered solution in Phosphate Buffer and NaCl (pH 6.3)
Sequence Length	152
Bioactivity	The ED50 as determined by the dose-dependent proliferation of Human TF-1 cells was found less than 0.5 ng/mL
Endotoxin Level	<1.0 EU/ug of recombinant protein as determined by the LAL method
Reconstitution	A quick spin of the vial followed by reconstitution in distilled water to a concentration not less than 0.1 mg/mL. This solution can then be diluted into other buffers.
Preparation and Storage	The lyophilized protein is stable for at least one year from date of receipt at -70 degree C. Upon reconstitution, this cytokine can be stored in working aliquots at 2-8 degree C for one month, or at -20 degree C for six months, with a carrier protein without detectable loss of activity. Avoid repeated freeze/thaw cycles.

IMAGES



Application Data



Application Data

ADDITIONAL INFORMATION

Related Product Information for IL3 active protein	IL3 is produced mainly by T cells following cell activation by antigens and mitogens, but also by keratinocytes, natural killer cells, mast cells, endothelial cells, and monocytes. The analysis of bacterial-derived recombinant IL3 shows that glycosylation is not required for activity. IL3 sequences are evolutionarily less well conserved with human and murine IL3 sharing approximately 29% homology (at the protein level) and murine and rat IL3 sharing approximately 54% homology. IL3 receptors are expressed on macrophages, mast cells, eosinophils, megakaryocytes, basophils, bone marrow progenitor cells and various myeloid leukemia cells. Binding of IL3 to its receptor causes specific phosphorylation of a 150 kDa membrane glycoprotein. Recombinant human IL3 is a non-glycosylated globular protein.
--	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	3002475
NCBI GeneID	3562
UniProt Accession #	P08700
Molecular Weight	15.0 kDa
NCBI Official Full Name	IL-3
NCBI Official Synonym Full Names	interleukin 3
NCBI Official Symbol	IL3
NCBI Official Synonym Symbols	IL-3; MCGF; MULTI-CSF
NCBI Protein Information	interleukin-3
UniProt Protein Name	Interleukin-3
UniProt Gene Name	IL3
UniProt Synonym Gene Names	IL-3; MCGF
UniProt Entry Name	IL3_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/e-coli-active-protein/il3/214236>