

MCP-2 (CCL8) Active Protein | MCP-2 active protein

Recombinant Human MCP-2 (CCL8)

CATALOG #
AAA214243

HOST
E Coli


PRODUCT SYNONYMS

MCP-2 (CCL8); Recombinant Human MCP-2 (CCL8); Monocyte Chemotactic Protein-2; MCP2; HC14; C-C motif chemokine 8; Recombinant Human Monocyte Chemoattractant Protein-2 (CCL8); MCP-2 active protein


PRODUCT SPECIFICATIONS

Host	E Coli
Purity/Purification	>95% as determined by SDS-PAGE
Form/Format	Lyophilized from a 0.2 um filtered 20 mM solution in PBS
Sequence Length	99
Bioactivity	The ED(50) was determined by the dose-dependent proliferation of Jurkat cells and was found to be <0.05ng/mL.
Endotoxin Level	<1.0 EU/ug of recombinant protein as determined by the LAL method
Reconstitution	A quick spin of the vial followed by reconstitution in distilled water to a concentration not less than 0.1 mg/mL. This solution can then be diluted into other buffers.
Preparation and Storage	The lyophilized protein is stable for at least one year from date of receipt at -70 degree C. Upon reconstitution, this cytokine can be stored in working aliquots at 2-8 degree C for one month, or at -20 degree C for six months, with a carrier protein without detectable loss of activity. Avoid repeated freeze/thaw cycles.

IMAGES



Application Data



Application Data

ADDITIONAL INFORMATION

Related Product Information for MCP-2 active protein	Monocyte chemotactic protein MCP-2 is a C-C chemokine. It shares over 60% amino acid identity with MCP-1 and MCP-3 and has about 30% identity with other C-C chemokines MIP-1 alpha, RANTES, and MIP-1 beta. MCP-2 is chemotactic for and activates a wide variety of inflammatory cells, and its spectrum of action on leukocytes is similar to that of MCP-3, including monocytes, T lymphocytes, natural killer cells, basophils, mast cells, and eosinophils, but differs from MCP-1, which is not active on eosinophils. MCP-2 has been proposed to interact with multiple C-C chemokine receptors, including those used by MCP-1 and MCP-3. However, since leukocytes express a multiplicity of chemokine receptors with promiscuous binding and functional properties, it is difficult to identify the receptors used by a given ligand on these cells.
--	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	2506280
NCBI GeneID	6355
UniProt Accession #	P80075
Molecular Weight	9.0 kDa
NCBI Official Full Name	C-C motif chemokine 8
NCBI Official Synonym Full Names	C-C motif chemokine ligand 8
NCBI Official Symbol	CCL8
NCBI Official Synonym Symbols	HC14; MCP2; MCP-2; SCYA8; SCYA10
NCBI Protein Information	C-C motif chemokine 8
UniProt Protein Name	C-C motif chemokine 8
UniProt Gene Name	CCL8
UniProt Synonym Gene Names	MCP2; SCYA10; SCYA8; MCP-2
UniProt Entry Name	CCL8_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/e-coli-active-protein/mcp-2-ccl8/214243>