

IFNA2 active protein

Recombinant Human IFNA2

CATALOG #
AAA214269

HOST
E Coli


PRODUCT SYNONYMS

IFNA2; Recombinant Human IFNA2; IFNA2A; IFN-alpha-2a; LeIF A; Recombinant Human Interferon-Alpha 2A (IFNA2); IFNA2 active protein


PRODUCT SPECIFICATIONS

Host	E Coli
Purity/Purification	>95% as determined by SDS-PAGE
Form/Format	Lyophilized from a 0.2 um filtered PBS solution pH 7.0.
Sequence Length	188
Bioactivity	The ED(50) was determined by the dose-dependent anti-proliferation of Ovcar3 cells and was found to be <20ng/mL.
Endotoxin Level	<1.0 EU/ug of recombinant protein as determined by the LAL method.
Reconstitution	A quick spin of the vial followed by reconstitution in distilled water to a concentration not less than 0.1 mg/mL. This solution can then be diluted into other buffers.
Preparation and Storage	The lyophilized protein is stable for at least one year from date of receipt at -70 degree C. Upon reconstitution, this cytokine can be stored in working aliquots at 2-8 degree C for one month, or at -20 degree C for six months, with a carrier protein without detectable loss of activity. Avoid repeated freeze/thaw cycles.

IMAGES



Application Data



Application Data

ADDITIONAL INFORMATION

Related Product Information for IFNA2 active protein	All known subtypes of IFN-alpha show the same antiviral antiparasitic, antiproliferative activities. Human IFN-alpha is also a potent antiviral substance in murine, porcine, and bovine cell systems. IFN-alpha forms are produced by monocytes/macrophages, lymphoblastoid cells, fibroblasts, and a number of different cell types following induction by viruses, nucleic acids, glucocorticoid hormones, and low-molecular weight substances. All IFN-alpha subtypes possess a common conserved sequence region between amino acid positions 115-151 while the amino-terminal ends are variable. Many IFN-alpha subtypes differ in their sequences at only one or two positions. IFN-alpha and IFN-beta are thought to bind to the same receptor. Signal transduction mechanisms elicited after binding of IFN-alpha to its receptors involves tyrosine phosphorylation (see also: PTK; protein tyrosine kinase) of various non-receptor tyrosine kinases belonging to the Janus kinases.
--	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	11067751
NCBI GeneID	3440
NCBI Accession #	NP_000596.2
NCBI GenBank Nucleotide #	NP_000596.2
UniProt Accession #	P01563
Molecular Weight	19.0 kDa
NCBI Official Full Name	interferon alpha-2
NCBI Official Synonym Full Names	interferon alpha 2
NCBI Official Symbol	IFNA2
NCBI Official Synonym Symbols	IFNA; INFA2; IFNA2B; IFN-alphaA
NCBI Protein Information	interferon alpha-2
UniProt Protein Name	Interferon alpha-2
UniProt Gene Name	IFNA2
UniProt Synonym Gene Names	IFNA2A; IFNA2B; IFNA2C; IFN-alpha-2; IeIF A
UniProt Entry Name	IFNA2_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/e-coli-active-protein/ifna2/214269>