

TNF active protein

Recombinant Mouse TNF

CATALOG #
AAA214321

HOST
E Coli


PRODUCT SYNONYMS

TNF; Recombinant Mouse TNF; TNF-alpha; TNF-a; Tumor necrosis factor ligand superfamily member 2; Cachectin; Tnfa; Tnfsf2; Tnf; Recombinant Mouse Tumor Necrosis Factor Alpha (TNF); TNF active protein


PRODUCT SPECIFICATIONS

Host	E Coli
Purity/Purification	>95% as determined by SDS-PAGE
Form/Format	Lyophilized from 0.2 um filtered PBS solution, pH 7.0.
Sequence Length	219
Bioactivity	The ED(50) was determined by cytolysis of murine L929 cells in the presence of Actinomycin D is ≤ 0.05 ng/mL, corresponding to a specific activity of $\geq 3.0 \times 10^7$ units/mg.
Endotoxin Level	<1.0 EU/ug of recombinant protein as determined by the LAL method.
Reconstitution	A quick spin of the vial followed by reconstitution in distilled water to a concentration not less than 0.1 mg/mL. This solution can then be diluted into other buffers.
Preparation and Storage	The lyophilized protein is stable for at least one year from date of receipt at -70 degree C. Upon reconstitution, this cytokine can be stored in working aliquots at 2-8 degree C for one month, or at -20 degree C for six months, with a carrier protein without detectable loss of activity. Avoid repeated freeze/thaw cycles.

IMAGES



Application Data



Application Data

ADDITIONAL INFORMATION

Related Product Information for TNF active protein	TNF is secreted by macrophages, monocytes, neutrophils, T cells, natural killer cells following their stimulation by bacterial lipopolysaccharides. Cells expressing CD4 secrete TNF-alpha while CD8(+) cells secrete little or no TNF-alpha. Stimulated peripheral neutrophilic granulocytes but also unstimulated cells and also a number of transformed cell lines, astrocytes, microglial cells, smooth muscle cells, and fibroblasts also secrete TNF. Human milk also contains this factor. The synthesis of TNF-alpha is induced by many different stimuli including interferons, IL-2, GM-CSF, SP, Bradykinin, immune complexes, inhibitors of cyclooxygenase and PAF (platelet activating factor). Human TNF-alpha is a non-glycosylated protein of 17 kDa and a length of 157 amino acids. Murine TNF-alpha is N-glycosylated. Homology with TNF-beta is approximately 30%. TNF-alpha forms dimers and trimers.
--	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	518831589
NCBI GeneID	21926
NCBI Accession #	NP_001265530.1
NCBI GenBank Nucleotide #	NP_001265530.1
UniProt Accession #	P06804
Molecular Weight	17.0 kDa
NCBI Official Full Name	tumor necrosis factor isoform 2
NCBI Official Synonym Full Names	tumor necrosis factor
NCBI Official Symbol	Tnf
NCBI Official Synonym Symbols	DIF; Tnfa; TNF-a; TNFSF2; Tnlg1f; Tnfsf1a; TNFalpha; TNF-alpha
NCBI Protein Information	tumor necrosis factor
UniProt Protein Name	Tumor necrosis factor
UniProt Gene Name	Tnf
UniProt Synonym Gene Names	Tnfa; Tnfsf2; TNF-a; NTF; ICD1; ICD2
UniProt Entry Name	TNFA_MOUSE

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/e-coli-active-protein/tnf/214321>