

Mouse gp100 / Melanosome / PMEL17 / SILV Monoclonal Antibody | anti-SILV antibody

gp100 / Melanosome / PMEL17 / SILV (Melanoma Marker) Mouse Monoclonal Antibody

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA214406	Mouse	Human.Does not react with Dog and Rat.Others not tested	Immunohistochemistry, Immunofluorescence, Flow Cytometry, Functional Assay


PRODUCT SYNONYMS

gp100 / Melanosome / PMEL17 / SILV; Monoclonal Antibody; gp100 / Melanosome / PMEL17 / SILV (Melanoma Marker) Mouse Monoclonal Antibody; 95kDa melanocyte-specific secreted glycoprotein; M-beta; Melanocyte lineage specific antigen GP100; Melanocyte protein Pmel 17; Melanoma associated ME20 antigen; Melanosomal matrix protein17; p100; p26; PMEL17; Premelanosome protein; Secreted melanoma-associated ME20 antigen; SILV; Silver homolog; anti-SILV antibody


PRODUCT SPECIFICATIONS

Host	Mouse
Reactivity	Human. Does not react with Dog and Rat. Others not tested
Clonality	Monoclonal
Isotype	IgG1, kappa
Clone Number	HMB45
Specificity	By immunohistochemistry, it specifically recognizes a protein in melanocytes and melanomas. This MAb reacts with junctional and blue nevus cells and variably with fetal and neonatal melanocytes. Intradermal nevi, normal adult melanocytes, and non-melanocytic cells are negative. It does not stain tumor cells of epithelial, lymphoid, glial, or mesenchymal origin. Metastatic amelanotic melanoma can often be confused with a variety of poorly differentiated carcinomas, large cell lymphomas, and sarcomas using H & E stains alone. It is also difficult to differentiate melanoma from spindle cell carcinomas and various types of mesenchymal neoplasms. Anti-HMB-45 stains fetal and neonatal melanocytes, junctional and blue nevus cells, and malignant melanoma. This MAb also stains Angiomyolipoma (PEComa).
Form/Format	Bioreactor Concentrate with 0.05% Azide.
Sequence Length	575
Applicable Applications for anti-SILV antibody	IHC (Immunohistochemistry), IF (Immunofluorescence), FCM/FACS (Flow Cytometry)
Cellular Localization	Cytoplasmic
Immunogen	Extract of pigmented melanoma metastases from lymph nodes
Hu-Chromosome Location	12q13.2
Positive Control	SK-MEL-28 cells or Melanoma
Preparation and Storage	Store at 2 to 8 degree C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

IMAGES



IHC (Immunohistochemistry)
Formalin-fixed, paraffin-embedded human Melanoma stained with gp100 / Melanosome Monoclonal Antibody (HMB45).



IHC (Immunohistochemistry)
Formalin-fixed, paraffin-embedded human Melanoma stained with gp100 / Melanosome Monoclonal Antibody (HMB45).

ADDITIONAL INFORMATION

References	Esclamado RM, et. al. American Journal of Surgery, 1986, 152(4):376-85. Gown AM, et. al. American Journal of Pathology, 1986, 123(2):195-203
------------	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	318037593
NCBI GeneID	6490
NCBI Accession #	NP_001186982.1
NCBI GenBank Nucleotide #	NM_001200053.1
UniProt Accession #	P40967; Q12763; Q14448; Q14817; Q16565; B3KS57; B7Z6D7
Molecular Weight	90-100kDa
NCBI Official Full Name	melanocyte protein PMEL isoform 2
NCBI Official Synonym Full Names	premelanosome protein
NCBI Official Symbol	PMEL
NCBI Official Synonym Symbols	P1; SI; SIL; ME20; P100; SILV; ME20M; gp100; ME20-M; PMEL17; D12S53E
NCBI Protein Information	melanocyte protein PMEL
UniProt Protein Name	Melanocyte protein PMEL
UniProt Gene Name	PMEL
UniProt Synonym Gene Names	D12S53E; PMEL17; SILV; ME20M; ME20-S; ME20S
UniProt Entry Name	PMEL_HUMAN

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/monoclonal/gp100-melanosome-pmel17-silv-antibody-clone-hmb45-immunohistochemistry-immunofluorescence-flow-cytometry-elisa/214406>