

Mouse Ep-CAM/CD326 Monoclonal Antibody | anti-Ep-CAM antibody

Ep-CAM/CD326 (Epithelial Marker)

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA214436	Mouse	Human.Does not react with Rat and Ferret. Others not known.	Immunohistochemistry, Western Blot, Immunofluorescence, Flow Cytometry, Functional Assay


PRODUCT SYNONYMS

Ep-CAM/CD326; Monoclonal Antibody; Ep-CAM/CD326 (Epithelial Marker); Adenocarcinoma-associated Antigen; Cell Surface Glycoprotein Trop-1; EGP2; EGP314; EGP40; Epithelial Cell Adhesion Molecule; Epithelial Glycoprotein 314; ESA; KSA; TACD1; TROP1; Tumor-associated Calcium Signal Transducer 1 (TACSTD1); ECS-1; Epidermal Surface Antigen 1; ESA1; FLOT2; Flotillin-2; Membrane Component; Chromosome 17; Surface Marker-1 (M17S1); REG-1; Reggie-1; Reggie-2; anti-Ep-CAM antibody


PRODUCT SPECIFICATIONS

Host	Mouse
Reactivity	Human. Does not react with Rat and Ferret. Others not known.
Clonality	Monoclonal
Isotype	IgG1, kappa
Clone Number	rVU-1D9
Specificity	This antibody reacts with the first EGF repeat in the extracellular domain of Ep-CAM. It is a 40-43kDa transmembrane epithelial glycoprotein, also identified as epithelial specific antigen (ESA), or epithelial cellular adhesion molecule (Ep-CAM). It is expressed on baso-lateral cell surface in most simple epithelia and a vast majority of carcinomas with the exception of adult squamous epithelium, hepatocytes and gastric epithelial cells. This antibody has been used to distinguish adenocarcinoma from pleural mesothelioma and hepatocellular carcinoma. This antibody is also useful in distinguishing serous carcinomas of the ovary from mesothelioma.
Form/Format	200ug/ml of recombinant MAb purified Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.
Sequence Length	314
Applicable Applications for anti-Ep-CAM antibody	IHC (Immunohistochemistry), WB (Western Blot), IF (Immunofluorescence), FCM/FACS (Flow Cytometry)
Immunogen	Small cell lung carcinoma cells
Cellular Localization	Cell Surface
Hu-chromosome Location	2p21
Positive Control	HT29 cells or breast tumor
Preparation and Storage	Antibody with azide - store at 2 to 8 degree C. Antibody without azide - store at -20 to -80 degree C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

IMAGES



IHC (Immunohistochemistry)
Formalin-fixed, paraffin-embedded human Thyroid Carcinoma stained with Cytokeratin 18 Monoclonal Antibody (Cocktail).



IHC (Immunohistochemistry)
Formalin-fixed, paraffin-embedded human Colon Carcinoma stained with Ep-CAM Recombinant Mouse Monoclonal Antibody (rVU-1D9).

ADDITIONAL INFORMATION

References	Tsubura A et. al. J Cut Pathol, 1992, 19:73-9. 1. Litvinov SV et. al. J Cell Biol, 1994, 125:437-46.
------------	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	218505670
NCBI GeneID	4072
NCBI Accession #	NP_002345.2
NCBI GenBank Nucleotide #	NM_002354.2
UniProt Accession #	P16422; P18180; Q6FG26; Q6FG49; Q96C47; Q9UCD0
Molecular Weight	40-43kDa
NCBI Official Full Name	epithelial cell adhesion molecule
NCBI Official Synonym Full Names	epithelial cell adhesion molecule
NCBI Official Symbol	EPCAM
NCBI Official Synonym Symbols	ESA; KSA; M4S1; MK-1; DIAR5; EGP-2; EGP40; KS1/4; MIC18; TROP1; EGP314; HNPCC8; TACSTD1
NCBI Protein Information	epithelial cell adhesion molecule
UniProt Protein Name	Epithelial cell adhesion molecule
UniProt Gene Name	EPCAM
UniProt Synonym Gene Names	GA733-2; M1S2; M4S1; MIC18; TACSTD1; TROP1; Ep-CAM; EGP; EGP314; hEGP314

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/monoclonal/ep-cam-cd326-antibody-clone-rvu-1d9-immunohistochemistry-western-blot-immunofluorescence-flow-cytometry-elisa/214436>