

Mouse Ep-CAM/CD326 Monoclonal Antibody | anti-Ep-CAM antibody

Ep-CAM/CD326 (Epithelial Marker)



CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA214438	Mouse	Human, Mouse, Rat. Others not known.	Immunohistochemistry

PRODUCT SYNONYMS

Ep-CAM/CD326; Monoclonal Antibody; Ep-CAM/CD326 (Epithelial Marker); Adenocarcinoma-associated Antigen; Cell Surface Glycoprotein Trop-1; EGP2; EGP314; EGP40; Epithelial Cell Adhesion Molecule; Epithelial Glycoprotein 314; ESA; KSA; TACD1; TROP1; Tumor-associated Calcium Signal Transducer 1 (TACSTD1); ECS-1; Epidermal Surface Antigen 1; ESA1; FLOT2; Flotillin-2; Membrane Component; Chromosome 17; Surface Marker-1 (M17S1); REG-1; Reggie-1; Reggie-2; anti-Ep-CAM antibody

PRODUCT SPECIFICATIONS

Host	Mouse
Reactivity	Human, Mouse, Rat. Others not known.
Clonality	Monoclonal
Isotype	IgG1
Clone Number	rEGP40/1110
Specificity	EGP40 is a 40-43kDa transmembrane epithelial glycoprotein, also identified as epithelial specific antigen (ESA), or epithelial cellular adhesion molecule (Ep-CAM). It is expressed on baso-lateral cell surface in most simple epithelia and a vast majority of carcinomas. This antibody has been used to distinguish adenocarcinoma from pleural mesothelioma and hepatocellular carcinoma. This antibody is also useful in distinguishing serous carcinomas of the ovary from mesothelioma.
Form/Format	200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.
Sequence Length	314
Applicable Applications for anti-Ep-CAM antibody	IHC (Immunohistochemistry)
Immunogen	Recombinant cytoplasmic-fragment of human EpCAM protein (around aa 280-350)
Cellular Localization	Cell Surface
Hu-chromosome Location	2p21
Positive Control	HT29 cells or Ovarian carcinoma
Preparation and Storage	Antibody with azide - store at 2 to 8 degree C. Antibody without azide - store at -20 to -80 degree C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.



IHC (Immunohistochemistry)
Formalin-fixed, paraffin-embedded Human Ovarian Carcinoma stained with EpCAM Mouse Recombinant Monoclonal Antibody (rEGP40/1110).

SDS-PAGE
SDS-PAGE Analysis Purified EpCAM Mouse Recombinant Monoclonal Antibody (rEGP40/1110).

ADDITIONAL INFORMATION

References	Tsubura A et. al. J Cut Pathol, 1992, 19:73-9. Litvinov SV et. al. J Cell Biol, 1994, 125:437-46.
------------	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	218505670
NCBI GeneID	4072
NCBI Accession #	NP_002345.2
NCBI GenBank Nucleotide #	NM_002354.2
UniProt Accession #	P16422; P18180; Q6FG26; Q6FG49; Q96C47; Q9UCD0
Molecular Weight	40-43kDa
NCBI Official Full Name	epithelial cell adhesion molecule
NCBI Official Synonym Full Names	epithelial cell adhesion molecule
NCBI Official Symbol	EPCAM
NCBI Official Synonym Symbols	ESA; KSA; M4S1; MK-1; DIAR5; EGP-2; EGP40; KS1/4; MIC18; TROP1; EGP314; HNPCC8; TACSTD1
NCBI Protein Information	epithelial cell adhesion molecule
UniProt Protein Name	Epithelial cell adhesion molecule
UniProt Gene Name	EPCAM
UniProt Synonym Gene Names	GA733-2; M1S2; M4S1; MIC18; TACSTD1; TROP1; Ep-CAM; EGP; EGP314; hEGP314

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.