

Rabbit Ep-CAM/CD326 Polyclonal Antibody | anti-Ep-CAM antibody

Ep-CAM/CD326 (Epithelial Marker)

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA214441	Rabbit	Human.Does not react with Mouse and Rat. Others not known.	Immunohistochemistry, Western Blot, Immunofluorescence, Flow Cytometry, Functional Assay


PRODUCT SYNONYMS

Ep-CAM/CD326; Polyclonal Antibody; Ep-CAM/CD326 (Epithelial Marker); Adenocarcinoma-associated Antigen; Cell Surface Glycoprotein Trop-1; EGP2; EGP314; EGP40; Epithelial Cell Adhesion Molecule; Epithelial Glycoprotein 314; ESA; KSA; TACD1; TACSTD1; TROP1; Tumor-associated Calcium Signal Transducer 1; ECS-1; Epidermal Surface Antigen 1; ESA1; FLOT2; Flotillin-2; M17S1 (Membrane Component; Chromosome 17; Surface Marker-1); REG-1; Reggie-1; Reggie-2; anti-Ep-CAM antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human. Does not react with Mouse and Rat. Others not known.
Clonality	Polyclonal
Isotype	IgG, kappa
Specificity	EGP40 is a 40-43kDa transmembrane epithelial glycoprotein, also identified as epithelial specific antigen (ESA), or epithelial cellular adhesion molecule (Ep-CAM). It is expressed on baso-lateral cell surface in most simple epithelia and a vast majority of carcinomas with the exception of adult squamous epithelium, hepatocytes and gastric epithelial cells. Ep-CAM antibody has been used to distinguish adenocarcinoma from pleural mesothelioma and hepatocellular carcinoma. Ep-CAM antibody is also useful in distinguishing serous carcinomas of the ovary from mesothelioma.
Form/Format	200ug/ml of Ab purified by Protein A. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.
Sequence Length	314
Applicable Applications for anti-Ep-CAM antibody	IHC (Immunohistochemistry), WB (Western Blot), IF (Immunofluorescence), FCM/FACS (Flow Cytometry)
Immunogen	Recombinant human TACSTD1 protein
Cellular Localization	Cell Surface
Hu-chromosome Location	2p21
Positive Control	HT29 cells or breast tumor
Preparation and Storage	Antibody with azide - store at 2 to 8 degree C. Antibody without azide - store at -20 to -80 degree C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

IMAGES



IHC (Immunohistochemistry)
Formalin-fixed, paraffin-embedded human Renal Cell Carcinoma stained with Ep-CAM Rabbit Polyclonal Antibody.



IHC (Immunohistochemistry)
Formalin-fixed, paraffin-embedded human Renal Cell Carcinoma stained with Ep-CAM Rabbit Polyclonal Antibody.

ADDITIONAL INFORMATION

References	Bjork, P., Jonsson, U., Svedberg, H., Larsson, K., Lind, P., Dillner, J., Hedlund, G., Dohlsten, M. and Kalland, T. 1993. Isolation, partial characterization, and molecular cloning of a human colon adenocarcinoma cell-surface glyco-protein recognized by the C215 mouse monoclonal antibody. J. Biol. Chem. 268: 24232-24241.
------------	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	218505670
NCBI GeneID	4072
NCBI Accession #	NP_002345.2
NCBI GenBank Nucleotide #	NM_002354.2
UniProt Accession #	P16422; P18180; Q6FG26; Q6FG49; Q96C47; Q9UCD0
Molecular Weight	40-43kDa
NCBI Official Full Name	epithelial cell adhesion molecule
NCBI Official Synonym Full Names	epithelial cell adhesion molecule
NCBI Official Symbol	EPCAM
NCBI Official Synonym Symbols	ESA; KSA; M4S1; MK-1; DIAR5; EGP-2; EGP40; KS1/4; MIC18; TROP1; EGP314; HNPCC8; TACSTD1
NCBI Protein Information	epithelial cell adhesion molecule
UniProt Protein Name	Epithelial cell adhesion molecule
UniProt Gene Name	EPCAM
UniProt Synonym Gene Names	GA733-2; M1S2; M4S1; MIC18; TACSTD1; TROP1; Ep-CAM; EGP; EGP314; hEGP314

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.