

Rabbit MUC1/CA15-3/EMA/CD227 Monoclonal Antibody | anti-MUC1 antibody

MUC1/CA15-3/EMA/CD227 (Epithelial Marker)

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA214457	Rabbit	Human. Others not known.	Immunohistochemistry, Immunofluorescence, Flow Cytometry, Functional Assay



PRODUCT SYNONYMS

MUC1/CA15-3/EMA/CD227; Monoclonal Antibody; MUC1/CA15-3/EMA/CD227 (Epithelial Marker); Breast carcinoma-associated antigen DF3; CA15-3; Carcinoma-associated mucin Episialin; Epithelial Membrane Antigen; H23AG; KL-6; MAM6; MUC-1; MUC-1/SEC; MUC-1/X; MUC1-alpha; MUC1-beta; MUC1-CT; MUC1-NT; MUC1/ZD; Mucin 1 cell surface associated; Mucin-1 subunit beta; Peanut-reactive urinary mucin; PEM; PEMT; Polymorphic epithelial mucin; PUM; Tumor-associated epithelial membrane antigen; anti-MUC1 antibody

PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human. Others not known.
Clonality	Monoclonal
Isotype	IgG
Clone Number	MUC1/1887R
Specificity	This MAb reacts with MUC1, a large transmembrane glycoprotein expressed on the ductal surface of normal glandular epithelia. It is used as tracer agent in CA15.3 assays. The extracellular domain of MUC1 largely consists of a highly conserved, O-glycosylated 20 amino acids tandem repeat which can occur 30-100 times per molecule depending on the length of the allele involved. In the vast majority of human carcinomas this protein is up-regulated and poorly glycosylated and appears on the cell surface in a non-polarized fashion. The dominant epitope of this MAb involves both amino acids as well as sugar moieties. Its epitope is destroyed by desialylation i.e. treatment with Neuraminidase.
Form/Format	200ug/ml of recombinant MAb purified by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.
Sequence Length	264
Applicable Applications for anti-MUC1 antibody	IHC (Immunohistochemistry), IF (Immunofluorescence), FCM/FACS (Flow Cytometry)
Immunogen	Human milk-fat globule membranes (HMFGM)
Cellular Localization	Cytoplasmic and Cell Surface
Hu-chromosome Location	1q22
Positive Control	MCF-7 or MDA-231 cells. Breast, colon, ovarian, endometrial carcinoma.
Preparation and Storage	Antibody with azide - store at 2 to 8 degree C. Antibody without azide - store at -20 to -80 degree C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

IMAGES



IHC (Immunohistochemistry)
Formalin-fixed, paraffin-embedded human Breast Carcinoma stained with MUC1 Rabbit Recombinant Monoclonal Antibody (MUC1/1887R).

SDS-PAGE
SDS-PAGE Analysis of Purified MUC1 Rabbit Recombinant Monoclonal Antibody (MUC1/1887R).

ADDITIONAL INFORMATION

References	Zotter et. al. Virchows Arch A Pathol Anat Histopathol. 1985; 406(2):237-51. Hilkens et al. Eur J Obstet Gynecol Reprod Biol. 1985 Jun; 19(6):356-7. Hilkens et al. Int J Cancer. 1984 Aug 15; 34(2):197-206.
------------	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	67189007
NCBI GeneID	4582
NCBI Accession #	NP_001018016.1
NCBI GenBank Nucleotide #	NM_001018016.2
UniProt Accession #	P15941; P13931; P15942; P17626; A5YRV1; A6ZID9; A6ZIE0; B1AVQ8; B1AVR0; B6ECA1; E7ESE5; E7EUG9
Molecular Weight	265-400kDa
NCBI Official Full Name	mucin-1 isoform 2
NCBI Official Synonym Full Names	mucin 1, cell surface associated
NCBI Official Symbol	MUC1
NCBI Official Synonym Symbols	EMA; MCD; PEM; PUM; KL-6; MAM6; MCKD; PEMT; CD227; H23AG; MCKD1; MUC-1; ADMCKD; ADMCKD1; CA 15-3; MUC-1/X; MUC1/ZD; MUC-1/SEC
NCBI Protein Information	mucin-1
UniProt Protein Name	Mucin-1
UniProt Gene Name	MUC1
UniProt Synonym Gene Names	PUM; MUC-1; CA 15-3; KL-6; PUM; PEM; EMA; MUC1-NT; MUC1-alpha; MUC1-beta

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.