

Mouse Thrombomodulin/CD141 Monoclonal Antibody | anti-THBD antibody

Thrombomodulin/CD141 (Endothelial Cell Marker)

CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA214533	Mouse	Human. Others not known.	Immunohistochemistry, Immunofluorescence, Flow Cytometry, Functional Assay, ELISA


PRODUCT SYNONYMS

Thrombomodulin/CD141; Monoclonal Antibody; Thrombomodulin/CD141 (Endothelial Cell Marker); AHUS6; BDCA3; CD141; Fetomodulin; Thbd; THPH12; THRM; Thrombomodulin (TM); anti-THBD antibody

PRODUCT SPECIFICATIONS


Host	Mouse
Reactivity	Human. Others not known.
Clonality	Monoclonal
Isotype	IgG2a, kappa
Clone Number	THBD/1782
Specificity	It recognizes a protein of 75kDa, identified as Thrombomodulin. Thrombomodulin is a transmembrane glycoprotein with natural anticoagulant properties. It is normally expressed by a restricted number of cells, such as endothelial and mesothelial cells. In addition, synovial lining and syncytio-trophoblasts of placenta also express thrombomodulin. This protein is present in almost all of benign vascular tumors and majority of malignant vascular tumors (Kaposi sarcoma, angiosarcoma, and epithelioid hemangioendothelioma). Hence, anti-thrombomodulin serves as a sensitive marker for lymphatic endothelial cells and their tumors. Recently, thrombomodulin antibody has been used for mesothelial cells and malignant mesotheliomas.
Form/Format	200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.
Sequence Length	575
Applicable Applications for anti-THBD antibody	IHC (Immunohistochemistry), IF (Immunofluorescence), FCM/FACS (Flow Cytometry), ELISA
Immunogen	Recombinant human Thrombomodulin protein fragment (aa69-194)
Cellular Localization	Cell Surface & Cytoplasmic
Hu-chromosome Location	20p11.2
Positive Control	THP1 Cells. Placenta, Angiosarcoma, Bladder or Cervical Carcinoma
Preparation and Storage	Antibody with azide - store at 2 to 8 degree C. Antibody without azide - store at -20 to -80 degree C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

IMAGES



IHC (Immunohistochemistry)

Formalin-fixed, paraffin-embedded human Cervical Carcinoma stained with Thrombomodulin/CD141 Monoclonal Antibody (THBD/1782).



IHC (Immunohistochemistry)

Formalin-fixed, paraffin-embedded human Placenta stained with Thrombomodulin/CD141 Monoclonal Antibody (THBD/1782).

ADDITIONAL INFORMATION

References	Appleton MA, Attanoos RL, Jasani B. Thrombomodulin as a marker of vascular and lymphatic tumours. Histopathology. 1996 Aug; 29(2):153-7.
------------	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	4507483
NCBI GeneID	7056
NCBI Accession #	NP_000352.1
NCBI GenBank Nucleotide #	NM_000361.2
UniProt Accession #	P07204; Q8IV29; Q9UC32
Molecular Weight	~100kDa
NCBI Official Full Name	thrombomodulin
NCBI Official Synonym Full Names	thrombomodulin
NCBI Official Symbol	THBD
NCBI Official Synonym Symbols	TM; THRM; AHUS6; BDCA3; CD141; THPH12
NCBI Protein Information	thrombomodulin
UniProt Protein Name	Thrombomodulin
UniProt Gene Name	THBD
UniProt Synonym Gene Names	THRM; TM

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/monoclonal/thrombomodulin-cd141-antibody-clone-thbd-1782-immunohistochemistry-immunofluorescence-flow-cytometry-elisa/214533>