

Rabbit TTF-1/NKX2.1 Monoclonal Antibody | anti-TTF-1 antibody

TTF-1/NKX2.1 (Thyroid & Lung Epithelial Marker)



CATALOG #	HOST	REACTIVITY	APPLICATIONS
AAA214538	Rabbit	Human, Mouse, Rat. Others not known.	Immunohistochemistry, Immunofluorescence, Flow Cytometry, Functional Assay

PRODUCT SYNONYMS

TTF-1/NKX2.1; Monoclonal Antibody; TTF-1/NKX2.1 (Thyroid & Lung Epithelial Marker); BCH; Benign chorea; BHC; Homeobox protein Nkx2.1; NK2 homeobox 1; NKX2.1; NKX2A; TEBP; Thyroid nuclear factor-1; Thyroid specific enhancer binding protein; Thyroid transcription factor-1 (TTF-1); Tin man; TITF1; TTF-1; anti-TTF-1 antibody


PRODUCT SPECIFICATIONS

Host	Rabbit
Reactivity	Human, Mouse, Rat. Others not known.
Clonality	Monoclonal
Isotype	IgG
Clone Number	NX2.1/1855R
Specificity	Recognizes a protein of 40kDa, identified as Thyroid transcription factor-1 (TTF-1). TTF-1 is a member of the NKx2 family of homeodomain transcription factors. It is expressed in epithelial cells of the thyroid gland and the lung. Nuclei from liver, stomach, pancreas, small intestine, colon, kidney, breast, skin, testes, pituitary, prostate, and adrenal glands are unreactive. Anti-TTF-1 is useful in differentiating primary adenocarcinoma of the lung from metastatic carcinomas originating in the breast, mediastinal germ cell tumors, and malignant mesothelioma. It can also be used to differentiate small cell lung carcinoma from lymphoid infiltrates. Loss of TTF-1 expression in non-small cell lung carcinoma has been associated with aggressive behavior of such neoplasms. TTF-1 reactivity is also seen in thyroid malignancies.
Form/Format	200ug/ml of recombinant MAb purified by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.
Sequence Length	401
Applicable Applications for anti-TTF-1 antibody	IHC (Immunohistochemistry), IF (Immunofluorescence), FCM/FACS (Flow Cytometry)
Immunogen	Recombinant full-length TTF-1 protein
Cellular Localization	Nuclear
Hu-chromosome Location	14q13.3
Positive Control	MAD109, MLE-15, H441-4, or H345 cells. Normal thyroid or lung.
Preparation and Storage	Antibody with azide - store at 2 to 8 degree C. Antibody without azide - store at -20 to -80 degree C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.



IHC (Immunohistochemistry)
Formalin-fixed, paraffin-embedded human Thyroid stained with TTF-1 Rabbit Recombinant Monoclonal Antibody (NX2.1/1855R).

Application Data



IHC (Immunohistochemistry)
Formalin-fixed, paraffin-embedded human Lung Adenocarcinoma stained with TTF-1 Rabbit Recombinant Monoclonal Antibody (NX2.1/1855R).

ADDITIONAL INFORMATION

References	Wert, S.E., et al. 2002. Increased expression of TTF-1 in respiratory epithelial cells inhibits alveolarization and causes pulmonary inflammation. Dev. Biol. 242: 75-87.
------------	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	118766339
NCBI GeneID	7080
NCBI Accession #	NP_001073136.1
NCBI GenBank Nucleotide #	NM_001079668.2
UniProt Accession #	P43699; O14954; O14955; Q7KZF6; Q9BRJ8; D3DSA3
Molecular Weight	40kDa
NCBI Official Full Name	homeobox protein Nkx-2.1 isoform 1
NCBI Official Synonym Full Names	NK2 homeobox 1
NCBI Official Symbol	NKX2-1
NCBI Official Synonym Symbols	BCH; BHC; NK-2; TEBP; TTF1; NKX2A; NMTC1; T/EBP; TITF1; TTF-1; NKX2.1
NCBI Protein Information	homeobox protein Nkx-2.1
UniProt Protein Name	Homeobox protein Nkx-2.1
UniProt Gene Name	NKX2-1

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com

Online datasheet: <https://stage.aaabiotech.com/monoclonal/ttf-1-nkx21-antibody-clone-nx2-1-1855r-immunohistochemistry-immunofluorescence-flow-cytometry-elisa/214538>