

Mouse GATA-3 Monoclonal Antibody | anti-GATA-3 antibody

GATA-3 (Breast and Urothelial Marker)

CATALOG # AAA214746	HOST Mouse	REACTIVITY Human. Others not known.	APPLICATIONS Immunohistochemistry
-------------------------------	----------------------	---	---

PRODUCT SYNONYMS

GATA-3; Monoclonal Antibody; GATA-3 (Breast and Urothelial Marker); GATA3; GATA binding protein-3; GATA-binding factor 3; HDR; HDRS; Transacting T-cell-specific transcription factor GATA-3; anti-GATA-3 antibody

PRODUCT SPECIFICATIONS

Host	Mouse
Reactivity	Human. Others not known.
Clonality	Monoclonal
Isotype	IgG2b, kappa
Clone Number	GATA3/2446
Specificity	GATA-3 is a zinc finger transcription factor and plays an important role in promoting and directing cell proliferation, development, and differentiation in many tissues and cell types. GATA-3 expression is primarily seen in breast and urothelial carcinomas. Therefore, GATA3 antibody can be useful in the identification of unknown primary carcinoma when carcinomas of the breast or bladder are a possibility.
Purity/Purification	Purified Ab with BSA and Azide at 200ug/ml OR Purified Ab WITHOUT BSA and Azide at 1.0mg/ml
Form/Format	200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.
Sequence Length	444
Applicable Applications for anti-GATA-3 antibody	IHC (Immunohistochemistry)
Immunogen	Recombinant fragment of human GATA3 protein (around aa 357-436) (exact sequence is proprietary)
Positive Control	MCF-7 or T47D Cells. Breast or Bladder Carcinoma.
Cellular Localization	Nuclear
Chromosome Location	10p14
Preparation and Storage	Antibody with azide - store at 2 to 8 degree C. Antibody without azide - store at -20 to -80 degree C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

IMAGES



Application Data

Analysis of Protein Array containing more than 19,000 full-length human proteins using GATA-3 Mouse Monoclonal Antibody (GATA3/2446). Z- and S- Score: The Z-score represents the strength of a signal that a monoclonal antibody (MAb) (in combination with a fluorescently-tagged anti-IgG secondary antibody) produces when binding to a particular protein on the HuProtTM array. Z-scores are described in units of standard deviations (SD's) above the mean value of all signals generated on that array. If targets on HuProtTM are arranged in descending order of the Z-score, the S-score is the difference (also in units of SD's) between the Z-score. S-score therefore represents the relative target specificity of a MAb to its intended target. A MAb is considered to specific to its intended target, if the MAb has an S-score of at least 2.5. For example, if a MAb binds to protein X with a Z-score of 43 and to protein Y with a Z-score of 14, then the S-score for the binding of that MAb to protein X is equal to 29.



IHC (Immunohistochemistry)

Formalin-fixed, paraffin-embedded human Breast Carcinoma stained with GATA-3 Mouse Monoclonal Antibody (GATA3/2446).

ADDITIONAL INFORMATION

References	Higgins JP, et al. Am J Surg Pathol. 2007; 31:673-680.; Liu H, et al. Am J Clin Pathol. 2012; 138:57-64.
------------	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	50541959
NCBI GeneID	2625
NCBI Accession #	NP_001002295.1
NCBI GenBank Nucleotide #	NM_001002295.1
UniProt Accession #	P23771; Q5VWG7; Q5VWG8; Q96J16
Molecular Weight	50kDa
NCBI Official Full Name	trans-acting T-cell-specific transcription factor GATA-3 isoform 1
NCBI Official Synonym Full Names	GATA binding protein 3
NCBI Official Symbol	GATA3
NCBI Official Synonym Symbols	HDR; HDRS
NCBI Protein Information	trans-acting T-cell-specific transcription factor GATA-3
UniProt Protein Name	Trans-acting T-cell-specific transcription factor GATA-3
UniProt Gene Name	GATA3

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.