

Mouse Nucleophosmin Monoclonal Antibody | anti-NPM1 antibody

Nucleophosmin (Acute Myeloid Leukemia Marker)


CATALOG # AAA214790	HOST Mouse	REACTIVITY Human. Others not tested.	APPLICATIONS Immunohistochemistry, Western Blot, ELISA
-------------------------------	----------------------	--	--

PRODUCT SYNONYMS

Nucleophosmin; Monoclonal Antibody; Nucleophosmin (Acute Myeloid Leukemia Marker); NO38; NPM; NPM1; Nucleolar phosphoprotein B23; Nucleolar protein NO38; Nucleophosmin (nucleolar phosphoprotein B23 numatrin); Nucleophosmin/nucleoplasmin family member 1; Numatrin; anti-NPM1 antibody


PRODUCT SPECIFICATIONS

Host	Mouse
Reactivity	Human. Others not tested.
Clonality	Monoclonal
Isotype	IgG2b, kappa
Clone Number	NPM1/1902
Specificity	Recognizes a 33kDa glycoprotein, identified as Nucleophosmin (NPM). It is predominantly localized in the nucleus of cells in most tissues. NPM is involved in ribosomal assembly and rRNA transport. It is an abundant protein that is highly phosphorylated by Cdc2 kinase during mitosis. This phosphoprotein moves between the nucleus and the cytoplasm. It is thought to be involved in several processes including regulation of the ARF/p53 pathway. A number of genes are fusion partners, in particular the anaplastic lymphoma kinase gene on chromosome 2. Mutations in exon 12 affecting the C-terminus of the protein are associated with an aberrant cytoplasmic location. Mutations in this gene are associated with acute myeloid leukemia. The antibody may be a useful aid for classification of acute myeloid leukemia.
Purity/Purification	Purified Ab with BSA and Azide at 200ug/ml OR Purified Ab WITHOUT BSA and Azide at 1.0mg/ml
Form/Format	200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.
Sequence Length	265
Applicable Applications for anti-NPM1 antibody	IHC (Immunohistochemistry), WB (Western Blot), ELISA
Immunogen	Recombinant human NPM1 protein fragment (aa185-287) (exact sequence is proprietary)
Positive Control	MCF-7, HeLa or A431 Cells. Skin, Colon
Cellular Localization	Nuclear and Cytoplasmic
Chromosome Location	5q35.1
Preparation and Storage	Antibody with azide - store at 2 to 8 degree C. Antibody without azide - store at -20 to -80 degree C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.




Application Data


Analysis of Protein Array containing more than 19,000 full-length human proteins using Nucleophosmin Mouse Monoclonal Antibody (NPM1/1902). Z- and S- Score: The Z-score represents the strength of a signal that a monoclonal antibody (MAb) (in combination with a fluorescently-tagged anti-IgG secondary antibody) produces when binding to a particular protein on the HuProt™ array. Z-scores are described in units of standard deviations (SD's) above the mean value of all signals generated on that array. If targets on HuProt™ are arranged in descending order of the Z-score, the S-score is the difference (also in units of SD's) between the Z-score. S-score therefore represents the relative target specificity of a MAb to its intended target. A MAb is considered to be specific to its intended target, if the MAb has an S-score of at least 2.5. For example, if a MAb binds to protein X with a Z-score of 43 and to protein Y with a Z-score of 14, then the S-score for the binding of that MAb to protein X is equal to 29.




WB (Western Blot)
Western Blot Analysis of A431 cell lysate using Nucleophosmin Mouse Monoclonal Antibody (NPM1/1902).



WB (Western Blot)
Western Blot (1) MCF-7 and (2) A431 Cell lysates using Nucleophosmin Mouse Monoclonal Antibody (NPM1/1902).



IHC (Immunohistochemistry)
Formalin-fixed, paraffin-embedded Human Colon Carcinoma stained with Nucleophosmin Mouse Monoclonal Antibody (NPM1/1902).



IHC (Immunohistochemistry)
Formalin-fixed, paraffin-embedded Human Skin stained with Nucleophosmin Mouse Monoclonal Antibody (NPM1/1902).

ADDITIONAL INFORMATION

References	Nozawa, Y., et al. 1996. Expression of nucleophosmin/B23 in normal and neoplastic colorectal mucosa. J. Pathol. 178: 48-52.
------------	---

NCBI AND UNIPROT INFORMATION

NCBI GI #	83641870
-----------	----------

NCBI GeneID	4869
NCBI Accession #	NP_001032827.1
NCBI GenBank Nucleotide #	NM_001037738.2
UniProt Accession #	P06748; P08693; Q12826; Q13440; Q13441; Q14115; Q5EU94; Q5EU95; Q5EU96; A8K3N7; B5BU00; D3DQL6
Molecular Weight	33kDa
NCBI Official Full Name	nucleophosmin isoform 3
NCBI Official Synonym Full Names	nucleophosmin 1
NCBI Official Symbol	NPM1
NCBI Official Synonym Symbols	B23; NPM
NCBI Protein Information	nucleophosmin
UniProt Protein Name	Nucleophosmin
UniProt Gene Name	NPM1
UniProt Synonym Gene Names	NPM; NPM

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/monoclonal/nucleophosmin-antibody-clone-npm1-1902-immunohistochemistry-western-blot-elisa/214790>