

Mouse Cyclin D2 Monoclonal Antibody | anti-CCND2 antibody

Cyclin D2

CATALOG # AAA214845	HOST Mouse	REACTIVITY Human. Others not known.	APPLICATIONS Immunohistochemistry
-------------------------------	----------------------	---	---

PRODUCT SYNONYMS

Cyclin D2; Monoclonal Antibody; CCND 2; ccnd2; CCND2_HUMAN; CyclinD2; G1/S specific cyclin D2; G1/S-specific cyclin-D2; KIAK0002; MGC102758; MPPH3; anti-CCND2 antibody

PRODUCT SPECIFICATIONS

Host	Mouse
Reactivity	Human. Others not known.
Clonality	Monoclonal
Isotype	IgG, kappa
Clone Number	CCND2/2620
Specificity	Cyclins are a family of proteins that control how cells proceed through the multi-step cycle of cell division. Cyclin D2 helps to regulate a step in the cycle called the G1-S transition, in which the cell moves from the G1 phase, when cell growth occurs, to the S phase, when the cell's DNA is copied (replicated) in preparation for cell division. Cyclin D2's role in the cell division cycle makes it a key controller of the rate of cell growth and division (proliferation) in the body.
Purity/Purification	Purified Ab with BSA and Azide at 200ug/ml OR Purified Ab WITHOUT BSA and Azide at 1.0mg/ml
Form/Format	200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.
Sequence Length	209
Applicable Applications for anti-CCND2 antibody	IHC (Immunohistochemistry)
Immunogen	Recombinant full-length human Cyclin D2 (CCND2) protein
Positive Control	Cervical Carcinoma. Esophagus.
Cellular Localization	Nuclear and Cytoplasmic
Chromosome Location	12p13.32
Preparation and Storage	Antibody with azide - store at 2 to 8 degree C. Antibody without azide - store at -20 to -80 degree C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

IMAGES



Application Data

Analysis of Protein Array containing more than 19,000 full-length human proteins using Cyclin D2 Mouse Recombinant Monoclonal Antibody (CCND2/2620). Z- and S- Score: The Z-score represents the strength of a signal that a monoclonal antibody (MAb) (in combination with a fluorescently-tagged anti-IgG secondary antibody) produces when binding to a particular protein on the HuProt™ array. Z-scores are described in units of standard deviations (SD's) above the mean value of all signals generated on that array. If targets on HuProt™ are arranged in descending order of the Z-score, the S-score is the difference (also in units of SD's) between the Z-score. S-score therefore represents the relative target specificity of a MAb to its intended target. A MAb is considered to specific to its intended target, if the MAb has an S-score of at least 2.5. For example, if a MAb binds to protein X with a Z-score of 43 and to protein Y with a Z-score of 14, then the S-score for the binding of that MAb to protein X is equal to 29.



IHC (Immunohistochemistry)

Formalin-fixed, paraffin-embedded Human Thyroid stained with Cyclin D2 Mouse Monoclonal Antibody (CCND2/2620).



IHC (Immunohistochemistry)

Formalin-fixed, paraffin-embedded Human Testis stained with Cyclin D2 Mouse Monoclonal Antibody (CCND2/2620).

ADDITIONAL INFORMATION

References	Draetta, G. 1990. Trends Biol. Sci. 15: 378-383.
------------	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	4502617
NCBI GeneID	894
NCBI Accession #	NP_001750.1
NCBI GenBank Nucleotide #	NM_001759.3
UniProt Accession #	P30279; Q13955; Q5U035; A8K531
Molecular Weight	34kDa
NCBI Official Full Name	G1/S-specific cyclin-D2
NCBI Official Synonym Full Names	cyclin D2
NCBI Official Symbol	CCND2
NCBI Official Synonym Symbols	MPPH3; KIAK0002
NCBI Protein Information	G1/S-specific cyclin-D2
UniProt Protein Name	G1/S-specific cyclin-D2
UniProt Gene Name	CCND2

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.