

Mouse CD80 Monoclonal Antibody | anti-CD80 antibody

CD80 (B7-1)

CATALOG # AAA214851	HOST Mouse	REACTIVITY Human. Others not known.	APPLICATIONS Immunohistochemistry, ELISA
-------------------------------	----------------------	---	--

PRODUCT SYNONYMS

CD80; Monoclonal Antibody; CD80 (B7-1); Activation B7-1 antigen; B lymphocyte activation antigen B7; B7; B7-1; BB1; CD28 antigen ligand 1; CD28LG; CD28LG1; Costimulatory factor CD80; CTLA-4 counter-receptor B7.1; LAB7; T-lymphocyte activation antigen CD80; anti-CD80 antibody

PRODUCT SPECIFICATIONS

Host	Mouse
Reactivity	Human. Others not known.
Clonality	Monoclonal
Isotype	IgG2b, kappa
Clone Number	C80/2776
Specificity	T cell proliferation and lymphokine production are triggered by occupation of the TCR by antigen, followed by a costimulatory signal that is delivered by a ligand expressed on antigen presenting cells. The B7-related cell surface proteins CD80 (B7-1) and CD86 (B7-2) are expressed on antigen presenting cells bind the homologous T cell receptors CTLA-4 (cytotoxic T lymphocyte-associated protein-4) and CD28 and trigger costimulatory signals for optimal T cell activation. CTLA-4 shares 31% overall amino acid identity with CD28 and it has been proposed that CD28 and CTLA-4 are functionally redundant. SLAM is a novel receptor on T cells that, when engaged, potentiates T cell expansion in a CD28-independent manner. B7, also designated BB1, is another ligand or counter receptor for CD28 and CTLA-4 that is expressed on the antigen-presenting cell.
Purity/Purification	Purified Ab with BSA and Azide at 200ug/ml OR Purified Ab WITHOUT BSA and Azide at 1.0mg/ml
Form/Format	200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.
Sequence Length	256
Applicable Applications for anti-CD80 antibody	IHC (Immunohistochemistry), ELISA
Immunogen	Recombinant fragment of human CD80 protein (around aa 35-142) (exact sequence is proprietary)
Positive Control	Raji or Ramos Cells. Tonsil, Spleen or Thymus
Cellular Localization	Cell Surface
Chromosome Location	3q13.3-q21
Preparation and Storage	Antibody with azide - store at 2 to 8 degree C. Antibody without azide - store at -20 to -80 degree C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

IMAGES



Application Data

Analysis of Protein Array containing more than 19,000 full-length human proteins using CD80 Mouse Monoclonal Antibody (C80/2776). Z- and S- Score: The Z-score represents the strength of a signal that a monoclonal antibody (MAb) (in combination with a fluorescently-tagged anti-IgG secondary antibody) produces when binding to a particular protein on the HuProtTM array. Z-scores are described in units of standard deviations (SD's) above the mean value of all signals generated on that array. If targets on HuProtTM are arranged in descending order of the Z-score, the S-score is the difference (also in units of SD's) between the Z-score. S-score therefore represents the relative target specificity of a MAb to its intended target. A MAb is considered to specific to its intended target, if the MAb has an S-score of at least 2.5. For example, if a MAb binds to protein X with a Z-score of 43 and to protein Y with a Z-score of 14, then the S-score for the binding of that MAb to protein X is equal to 29.



IHC (Immunohistochemistry)

Formalin-fixed, paraffin-embedded human Tonsil stained with CD80 Mouse Monoclonal Antibody (C80/2776).

ADDITIONAL INFORMATION

References	Peach, R.J., et al. 1995. J. Biol. Chem. 270: 21181-21187.
------------	--

NCBI AND UNIPROT INFORMATION

NCBI GI #	4885123
NCBI GeneID	941
NCBI Accession #	NP_005182.1
NCBI GenBank Nucleotide #	NM_005191.3
UniProt Accession #	P33681; Q5DTA9; Q5DTB0
Molecular Weight	60kDa
NCBI Official Full Name	T-lymphocyte activation antigen CD80
NCBI Official Synonym Full Names	CD80 molecule
NCBI Official Symbol	CD80
NCBI Official Synonym Symbols	B7; BB1; B7-1; B7.1; LAB7; CD28LG; CD28LG1
NCBI Protein Information	T-lymphocyte activation antigen CD80
UniProt Protein Name	T-lymphocyte activation antigen CD80
UniProt Gene Name	CD80
UniProt Synonym Gene Names	CD28LG; CD28LG1; LAB7; B7

For Research Use Only. Not for use in diagnostic or therapeutic procedures. Small volumes of product may become entrapped in the seal of the vial during shipment and storage; if necessary, briefly centrifuge the vial on a tabletop centrifuge to dislodge any liquid in the cap.

AAA Biotech | Tel: +1 (619) 604-9440 | Fax: +1 (619) 604-9441 | contact@aaabiotech.com
Online datasheet: <https://stage.aaabiotech.com/monoclonal/cd80-antibody-clone-c80-2776-immunohistochemistry-elisa/214851>